

Original Research Articles

Effects of dietary β -sitosterol supplementation on growth performance, antioxidant ability, and disease resistance in largemouth bass *Micropterus salmoides*

Yangyang Xing^{1,2}, Liping Zhang³, Mingyang Xue¹, Wei Liu⁴, Nan Jiang⁴, Yiqun Li¹, Jianwu Chen¹, Yuding Fan¹, Yong Zhou¹, Yan Meng^{1a}

¹ Chinese Academy of Fishery Sciences (China), Yangtze River Fisheries Research Institute (China), ² Shanghai Ocean University (China), College of Fisheries and Life Science (China), ³ Chongqing Fisheries Technical Extension Center (China), Chongqing Fisheries Technical Extension Center (China), ⁴ Chinese Academy of Fishery Sciences (China), Yangtze River Fisheries Research Institute (China)

Keywords: Largemouth bass, β -sitosterol, growth performance, antioxidant ability, disease resistance https://doi.org/10.46989/001c.117345

Israeli Journal of Aquaculture - Bamidgeh

Vol. 76, Issue 2, 2024

 β -sitosterol, as the most abundant phytosterol, has been shown to exert multiple biological roles in in vitro and in vivo animal studies. Largemouth bass (Micropterus salmoides) is an economical freshwater-farmed species widely cultivated in China due to its nutritious and delicious meat. This study investigated the effects of dietary β -sitosterol supplementation on largemouth bass farming. In the study, largemouth bass were fed with five levels of β -sitosterol (0, 20, 40, 80, and 160 mg/kg) supplementation in a commercial feed for four weeks. The growth performance, antioxidant ability, intestinal structure, immune-related gene expression, and resistance to pathogens such as Aeromonas hydrophila and largemouth bass virus (LMBRaV) were detected. The results revealed that β -sitosterol supplementation in feed increased the weight gain rate of largemouth bass, with 40 mg/kg supplementation displaying the best effect. The serum biochemical indices all showed an increase, including glucose and alkaline phosphatase. The intestinal villus length and muscularis propria thickness also increased, accompanied by elevated digestive enzyme activities, which promoted digestion and absorption of nutrients. Meanwhile, the levels of total cholesterol, aspartate aminotransferase, and alanine aminotransferase in serum decreased, indicating reduced liver damage. In addition, β -sitosterol supplementation effectively enhanced the antioxidant capacity of largemouth bass by decreasing malondialdehyde and elevating superoxide dismutase. Immune-related gene expression also changed. Pathogen infection tests revealed that β -sitosterol supplementation in feed had a certain protective effect against infection in largemouth bass, with the protection rate against A. hydrophila being higher than LMBRaV. The 40 mg/kg β -sitosterol supplementation group exhibited the best results. Collectively, the results revealed that the β -sitosterol can promote growth, improve enzyme activity, stimulate intestinal digestion, increase immune gene expression, and induce disease resistance in largemouth bass, with the optimal supplementation dosage being 40 mg/kg. β -sitosterol might be an excellent feed additive in aquaculture.

INTRODUCTION

Largemouth bass (*Micropterus salmoides*) was introduced in China from America in 1983. It is a thriving highly economical freshwater-farmed species that has been widely cultivated in China due to its nutritious and delicious meat.¹ Under intensive aquaculture conditions, the growth rate of largemouth bass is very fast and susceptible to pathogens due to environmental stresses such as poor water quality and farm treatment. These diseases lead to severe economic losses in largemouth bass breeding.² The major pathogens in largemouth bass include bacterial pathogens such as *Nocardia* and *Aeromonas hydrophila* (*A. hydrophila*), viral pathogens such as Largemouth bass ranavirus (LMBRaV), and parasitic pathogens such as *Trichodina*, *Apiosoma blanchard*, and *Ichthyophthirius*.³ Therefore, many studies have investigated the pathogenicity, detection and diagnostic technologies, prevention and control measures, medicine, and vaccines to prevent these diseases.⁴ Antibiotics or

a Corresponding author. Yan Meng, E-mail address: mengy@yfi.ac.cn

drugs are often used to control diseases in aquaculture. However, these treatments face various challenges, including potential drug resistance, immune system suppression, environmental hazards, and food safety problems.⁵ Nonetheless, no effective and commercialized vaccine or drug has been developed for disease control of largemouth bass.⁶

Functional feed or feed additives isolated from natural herbs can enhance aquatic animal health and improve the water environment in aquaculture.⁷ The application of naturel-identical compounds in feed additives to stimulate the immune system of fish and prevent and control diseases has emerged as a research hotspot.⁶ Plant sterols (phytosterols) are naturally occurring plant constituents that are structurally similar to cholesterol. They are active components present in plants and have a variety of pharmacological activities, such as anti-inflammation, anti-diabetes, and anti-cancer properties. Natural plant-derived steroid compounds are novel feed additives that regulate immunity and growth in animals.⁷ Phytosterols mainly include βsitosterol, stigmasterol, rapeseed oil sterols, and rapeseed sterols. Among them, β -sitosterol is the most abundant and plays an essential role in physiological processes . β-sitosterol was found to exert significant anti-inflammatory effects in animals.⁸ β-sitosterol also resulted in several effects in mice, including increased body weight, improved colon morphology, and reduced expression of proinflammatory factors.⁹ It reduced Pseudomonas aeruginosa lung infection and inflammation in mice,¹⁰ and dramatically increased the expression of antimicrobial proteins in intestinal epithelial cells in mice.⁹ However, studies on the effects of β -sitosterol on fish are very limited. One piece of research reported that β -sitosterol enhanced the intestinal immune function of large yellow croakers (Larimichthys crocea) infected with A. hvdrophila.¹¹ Furthermore, B-sitosterol was found to exert certain anti-inflammatory and antioxidant effects in zebrafish inflammation induced by copper sulfate.¹² Therefore, the current study was designed to investigate the effects of β -sitosterol supplementation on growth performance, antioxidant ability, and disease resistance in largemouth bass. In addition, different doses were tested to determine the optimal dose. The results are expected to provide evidence facilitating the development of natural functional feed for largemouth bass, thereby improving its health for the sustainable development of the breeding industry.

MATERIALS AND METHODS

FISH AND EXPERIMENTAL DIETS

Largemouth bass were purchased from one farm of Wuhan, Hubei Province, China. All largemouth bass were kept in fifteen recirculating aquaculture systems and fed with commercial largemouth bass feed (Tongwei, China) for 14 days before the experiment. The largemouth bass (weight $37.30 \pm$ 0.26 g) were randomly assigned to five feeding treatments, each consisting of three replicates and 60 largemouth bass per replicate. β -sitosterol was purchased from Yuanye Bio-Technology Co., Ltd. (Shanghai, China), which was rolled into powder and mixed with flour before being added to commercial feed. Five different concentrations of β -sitosterol (0, 20, 40, 80, and 160 mg/kg) were added to the commercial feeds according to the previous study results.^{13,14} The prepared supplemented diet was labeled as control (0), T1 (20), T2 (40), T3 (80), and T4 (160). The largemouth bass were kept in oxygenated water at a temperature of 25 ± 1°C. The fish were fed with the prepared supplemented diet three times a day for 28 days until the end of the experiment. All the experimental procedures were approved by the Yangtze River Fisheries Research Institute's Animal Experimental Ethical Committee (YFI 2022-Mengyan-0412).

SAMPLE COLLECTION

Blood samples were collected for serum biochemical analysis, and tissue samples for histological analysis, digesting and antioxidant enzyme activity analysis, and gene expression analysis. After the feeding experiment, largemouth bass were anesthetized with MS-222 (100 mg/L, Sigma, USA) and weighed. Nine largemouth bass from each experimental group were randomly selected and blood was collected from the tail vein into 1.5 mL centrifugal tubes. The collected blood was kept at 4°C for 2 h and then centrifuged at 3000 rpm for 10 min. The supernatant was obtained and preserved at -80°C. For histological analysis, the tissues from the midgut and liver were soaked in a 4% paraformaldehyde solution. In addition, the liver and intestinal tissues were carefully removed and preserved at -80°C for the intestinal digesting and antioxidant enzyme activity assay. At days 3, 7, 14, 21, and 28 of the feed experiment, the head kidney and spleen tissues from three largemouth bass in each tank were collected and kept in TRIzol (Invitrogen, USA) to extract RNA for the gene expression analysis.

GROWTH PERFORMANCE

The growth performance of weight gain rate (WGR), hepatosomatic index (HSI), viscerosomatic index (VSI), condition factor (CF), and feed conversion ratio (FCR) were evaluated to assess the effects of dietary β -sitosterol supplementation on the growth performance of largemouth bass. The calculation formulae for these parameters were obtained from a previously published study¹⁵: WGR (%) = $100 \times (W_t - W_0)/W_0$; FCR (%) = $100 \times F/(W_t - W_0)$; CF (g/cm³) = $100 \times W_t/$ (body length)¹³; HSI (%) = $100 \times \text{liver}$ weight/ W_t ; VSI (%) = $100 \times \text{visceral weight}/W_t$. In these formulae, W_t and W_0 represent the final weight and initial weight of the largemouth bass, and the F indicates the total quantity of feed consumed.

SERUM BIOCHEMICAL ANALYSIS

In this study, the serum biochemical analysis indicators included glucose (GLU), total cholesterol (TCHO), triglyceride (TG), aspartate aminotransferase (AST), alanine aminotransferase (ALT), alkaline phosphatase (ALP), total protein (TP), and albumin (ALB). These parameters were measured using different commercial kits (Dibao, China) and an

Genes	Primer sequence	Reference/Accession no.	
β-actin	F:5'-CAGGATGCAGAAGGAGATCACA-3' R:5'-CTCCTGCTTGCTGATCCACAT-3'	Yi et al., ¹⁶	
IL-1ß	F:5'-TGGTGGAAAACAGCATGGAGC-3' R: 5'-AGGGTGCACGTAGTTCGACA-3'	Yi et al., ¹⁶	
TNF-α	F: 5′-ACTTCGTCTACAGCCAGGCA-3′ R: 5′-AGTAACGCGAGACCCTGTGG-3′	Yi et al., ¹⁶	
IL-10	F: 5'-TCAGGAGGTGGAGCAGGAC-3' R: 5'-AGGACCAAGATGGCAACG-3'	Nie et al., ¹⁷	
IRF3	F: 5'-CCACTGCTCATCCCATGGCT-3' R: 5'-GAGGAGTCCTGTCTTAAACC-3'	XM_038735465.1	
Мх	F: 5'-TAAAATGGCTGGGGTCGGGG-3' R: 5'-CATTGCACGGAACGACCACC-3'	Yi et al., ¹⁶	

Note: IL-1 β : interleukin-1 beta, TNF- α : tumor necrosis factor- α , IL-10: interleukin-10, IRF3: interferon regulatory factor 3, Mx: myxovirus resistance.

Olympus 600 automatic biochemical analyzer (Olympus, Japan).

INTESTINAL DIGESTIVE AND ANTIOXIDANT ENZYME ACTIVITIES ANALYSIS

The levels of trypsin, amylase, and lipase in the intestine, and superoxide dismutase (SOD), malondialdehyde (MDA), total cholesterol (TCHO), and total cholesterol (TCHO) in the liver were determined to assess the intestinal digestive and antioxidant enzyme activities of largemouth bass. They were detected by the corresponding detection test kits (Jiancheng, China), and analyzed using a UV spectrophotometer (Thermo, USA) and microplate reader (Biotek, USA).

IMMUNE-RELATED GENE EXPRESSION ANALYSIS

Total RNA was isolated from tissue samples collected from the head kidney and spleen, conserved in TRIzol reagent following the instructions. Subsequently, the total RNA was reverse-transcribed into cDNA using a reverse transcription kit (TaKaRa, China). Immune-related genes were selected based on previously published studies, including tumor necrosis factor- α (*TNF-\alpha*), myxovirus resistance (*Mx*) and interleukin-1 beta (*IL-1\beta*),¹⁶ interferon regulatory factor 3 (*IRF3*) (XM_038735465.1), and interleukin-10 (*IL-10*).¹⁷ In this study, β -actin was used as an internal reference gene.¹⁶ The primer sequences are listed in Table 1.

CHALLENGE TEST

Aeromonas hydrophila and largemouth bass ranavirus (LM-BRaV) were isolated and stored in our laboratory.³ Largemouth bass were infected to evaluate the effects of β -sitosterol on disease resistance. After the supplemented feeding experiment, thirty fish were chosen randomly and infected with 0.2 mL A. hydrophila suspension (1.0 × 10⁷ colony forming unit CFUs/mL) intraperitoneally. Another thirty fish were chosen randomly and infected with 0.1 mL LM-

BRaV of 1×10^3 TCID₅₀ via intraperitoneal injection. In addition, thirty fish were injected with phosphate-buffered saline to establish the mock group. The deaths in each group were recorded daily. To ensure that largemouth bass were killed by *A. hydrophila* and LMBRaV, the diseased fish were collected, and pathogens were detected.

RESULTS

EFFECTS OF DIETARY **B**-SITOSTEROL SUPPLEMENTATION ON GROWTH PERFORMANCE

A 100% survival rate was achieved in all experimental groups during the experiment period. All the growth parameters results of largemouth bass are displayed in Table 2. The final weight increased in all experimental groups, with the T2 group displaying the most significant increase compared to the control group (P < 0.05). Weight gain rate (WGR) and condition factor (CF) demonstrated a similar increase compared to the control group (P < 0.05). In contrast, a decreasing trend was observed for the feed conversion ratio (FCR), viscerosomatic index (VSI) and hepatosomatic index (HSI) across the β -sitosterol treatment groups (P < 0.05).

RESULTS OF SERUM BIOCHEMICAL ANALYSIS

Serum biochemical indices of β -sitosterol supplementation to largemouth bass feed were displayed in **Figure 1**. The glucose (GLU) levels were significantly increased in the β sitosterol treatment groups compared to the control group (P < 0.05). Moreover, the total cholesterol (TCHO) and alanine aminotransferase (ALT) were significantly decreased in β -sitosterol treatment groups compared to the control group (P < 0.05). The alkaline phosphatase (ALP), total protein (TP), and albumin (ALB) levels exhibited the same increase when compared to the control group.

Table 2. Effects of p-sitosterol supplementation on growth performance of largemouth	n bass
--	--------

Items	Groups					
	control	T1 (20)	T2 (40)	T3 (80)	T4 (160)	
Initial weight (g)	36.75 ± 0.22^{a}	38.17 ± 0.09^{b}	36.52 ± 0.16^{a}	36.83 ± 0.72^{a}	38.25 ± 0.10^{b}	
Final weight (g)	81.65 ± 0.88^{a}	88.73 ± 0.64 ^c	$90.21 \pm 0.61^{\circ}$	85.64 ± 0.88^{b}	85.44 ± 0.29^{b}	
WGR (%)	122.20 ± 3.01^{a}	132.48 ± 1.61 ^a	147.04 ± 1.79 ^b	132.73 ± 6.02 ^a	123.37±0.44 ^a	
VSI (%)	7.43 ± 0.13^{d}	$6.81 \pm 0.02^{\circ}$	$6.81 \pm 0.02^{\circ}$	6.29 ± 0.09 ^b	5.19 ± 0.06 ^a	
HSI (%)	$2.41 \pm 0.15^{\circ}$	2.33 ± 0.03^{bc}	2.08 ± 0.00 ^{ab}	2.05 ± 0.09^{a}	1.84 ± 0.06 ^a	
CF (g/cm ³)	3.25 ± 0.03^{a}	3.48 ± 0.02^{b}	$3.63 \pm 0.03^{\circ}$	3.31 ± 0.03^{a}	3.53 ± 0.01^{b}	
FCR (%)	1.11 ± 0.02^{d}	0.99 ± 0.01^{b}	0.93 ± 0.01^{a}	1.03 ± 0.03^{bc}	1.06 ± 0.00^{cd}	

Note: WGR: Weight gain rate, VSI: viscerosomatic index, HSI: hepatosomatic index, CF: condition factor, FCR:feed conversion ratio.





Figure 1. Effects of β -sitosterol supplementation on serum biochemical indices of largemouth bass.

GLU: glucose, TCHO: total cholesterol, TG: triglyceride, ALB: albumin, TP: total protein, ALT: alanine aminotransferase, AST: aspartate aminotransferase, ALP: alkaline phosphatase. Different superscripts within a row indicate significant differences (P < 0.05).

ANALYSIS OF INTESTINAL DIGESTIVE AND ANTIOXIDANT ENZYME ACTIVITIES

The results of intestinal digestive and antioxidant enzyme activities are displayed in Figure 2. The results indicated that β -sitosterol supplementation can lead to changes in intestinal digestive and antioxidant enzyme activity. As shown in Figure 2A, total cholesterol (TCHO) and malon-dialdehyde (MDA) were significantly decreased in β -sitosterol treatment groups compared to the control group (P < 0.05). Meanwhile, superoxide dismutase (SOD) was increased following β -sitosterol supplementation but showed no statistical significance (P > 0.05). There were no significant differences in serum triglyceride (TG) contents among the treatment groups (P > 0.05). However, trypsin, lipase, and amylase were significantly increased following β -sitosterol treatments compared to the control group. The most

significant increase was in the T2 group (P < 0.05) (Figure 2B).

HISTOPATHOLOGICAL ANALYSIS

Histopathological analysis indicated that all groups of liver tissues exhibited no abnormal changes. Compared to the control group, the β -sitosterol supplementation groups showed clear cell-to-cell boundaries, and no inflammatory infiltration or hepatocellular damage was observed (Figure <u>3A</u>). In terms of intestinal tissue structure, the β -sitosterol supplementation treatment groups exhibited normal microstructure and dense microvilli (Figure <u>3B</u>). β -sitosterol supplementation groups demonstrated increased intestinal villus length and muscularis propria thickness compared to the control group (Figure <u>3C</u>).





Figure 2. Effects of β -sitosterol supplementation on intestinal digestive and antioxidant enzyme activities of largemouth.

(A) Antioxidant enzyme activities; (B) intestinal digestive activities. MDA: malondialdehyde, SOD: superoxide dismutase, TCHO: total cholesterol, TG: triglyceride. Different superscripts within a row indicate significant differences (P < 0.05).







Figure 3. Analysis of liver morphology (A), intestinal morphology (B) and intestine histometric (C) of largemouth bass kept in the express for 28 days while fed meals with varying amounts of β -sitosterol supplementation.

Different superscripts within a row indicate significant differences (P < 0.05).

ANALYSIS OF THE EXPRESSION OF IMMUNE-RELATED GENES

The expression levels of immune-related genes (*IL-10*, *IL-1β*, *TNF-α*, *IRF-3*, and *Mx*) were measured in the largemouth bass's head kidney and spleen on days 3, 7, 14, 21, and 28. Decreased expression levels of *IL-1β* and *TNF-α* were observed in all β-sitosterol treatment groups, with the most significant reduction found in the head kidney on day 14 and in the spleen on day 21 (P < 0.05). The expression level of *IL-10* was elevated in all β-sitosterol treatment groups, with the highest in the head kidney on day 14 and in the spleen on day 21 (P < 0.05). *IRF3* and *Mx* expression levels were higher in all β-sitosterol treatment groups; the most significant increase was observed in the head kidney on day 3 and in the spleen on day 14 (**Figure 4**).

CHALLENGE TEST

To investigate the protective effect of sterol on largemouth bass, the fish were infected with LMBRaV and *A. hydrophila* to carry out the infection experiments. During the challenge tests, no death occurred in the mock group. In the



Figure 4. Effects of β -sitosterol supplementation on the expression of immune-related genes in the head kidney and spleen of largemouth bass.

Different superscripts within a row indicate significant differences (P < 0.05).

LMBRaV challenge test, largemouth bass began to die from day 3 in the control group (commercial feed without β sitosterol supplement) and from day 4 in the β -sitosterol treatment groups. The LMBRaV cumulative survival rate reached 20% in the T2 group (**Figure 5A**). In the *A. hydrophila* challenge test, all largemouth bass died on day 6 in the control group, whereas the survival rates for T1, T2, T3, and T4 groups were 30, 40, 40 and 35%, respectively (Figure 5B). These findings indicated that compared to the control group, the β -sitosterol treatment groups had higher protective efficiency than largemouth bass. β -sitos-



Figure 5. Cumulative survival rate of largemouth bass after LMBRaV (A) and *A. hydrophila* (B) challenge.

**P < 0.01; *P < 0.05.

terol supplementation displayed higher protection against *A. hydrophila* than LMBRaV.

DISCUSSION

With the development of aquaculture and aquafeed additives, fish diets supplemented with effective plant extracts can enhance growth performance, alter gut microbiota, improve immune and oxidative functions, and control bacterial infections, thereby enhancing healthy breeding.^{2,6} This study showed that adding β -sitosterol to the largemouth bass feed can improve the growth performance, antioxidant ability, intestinal structure, immune-relative gene expression, and disease resistance. The results not only expand the use of β -sitosterol but also provide a potentially functional plant-derived feed additive for the healthy development of the largemouth bass breeding industry.

Many studies have reported that phytosterols can improve animal growth performance. Adding 75 mg/kg βsitosterol to broiler feed significantly increased the body weight and feed intake.¹⁸ Supplementing phytosterols to the feed of weaned piglets could increase their intestinal villus length.¹⁹ In tilapia fish, the addition of 20 and 40 mg/kg phytosterols to feed significantly increased the final weight, weight gain rate (WGR), and specific growth rate and reduced the feed conversion ratio (FCR).²⁰ In the present study, the β -sitosterol supplement groups demonstrated superior weight gain rate (WGR) and condition factor (CF), and lower feed conversion ratio (FCR) compared to the control group (only commercial feed). Meanwhile, the T2, T3, and T4 groups exhibited significantly increased intestinal villus length and muscularis propria thickness of largemouth bass. Supplementation with 40 mg/kg showed the best results. The findings led to a similar conclusion to

previous studies, indicating that β -sitosterol exerts a positive effect on growth performance in animals.

Previous studies have also shown that phytosterols impact animal blood biochemical indexes and digestive enzyme activity, which is conducive to animal growth, digestion, and absorption. Digestive enzyme activity has a direct impact on the ability to absorb and digest nutrients, and is an essential measure of fish growth performance. Alkaline phosphatase (ALP) is closely related to the transfer of phosphate groups and calcium and phosphorus metabolism. β sitosterol can promote hepatic glycogenolysis.²¹ The addition of β -sitosterol to rat feed increased serum glucose (GLU).²² Furthermore, adding 40 mg/kg phytosterol to the basal diet of broiler chickens resulted in significantly elevated levels of trypsin, lipase, and amylase activities.²³ Phytosterols (particularly β -sitosterol) can enhance the synthesis of animal proteins. In this study, the changes in serum biochemical indices, intestinal morphology, and antioxidant enzyme activities when adding β -sitosterol in largemouth bass commercial feed were investigated. The results revealed increased serum biochemical indices, including glucose and alkaline phosphatase. The intestinal villus length and muscularis propria thickness increased too, accompanied by the digestive enzyme activities, which all can promote digestion and absorption of nutrients. Meanwhile, the levels of total cholesterol, aspartate aminotransferase, and alanine aminotransferase in serum decreased to reduce liver damage. In addition, β -sitosterol supplementation can enhance the antioxidant capacity of largemouth bass by decreasing malondialdehyde and elevating superoxide dismutase. In fish, antioxidant enzymes play an essential role in the antioxidant system. β -sitosterol provides protection against oxidative stress by regulating antioxidant enzymes and free radicals.²⁴ Aspartate aminotransferase (AST) and alanine aminotransferase (ALT) levels are often used as specific indicators of impaired liver function. Total cholesterol (TCHO) reflects the body's lipid metabolism. In addition, superoxide dismutase (SOD) activity showed an increasing trend in β -sitosterol treatment groups in tilapia.²⁰ β-sitosterol also reduced total cholesterol (TCHO) in the serum and liver. Collectively, these findings suggest that dietary β-sitosterol supplementation promoted the growth performance of largemouth bass by affecting the intestinal morphology and serum biochemical factors, increasing the hepatic antioxidant capacity to improve the absorption of nutrients.

Abundant evidence suggests that certain pro-inflammatory cytokines (like *TNF-* α and *IL-1* β) and anti-inflammatory cytokines (like *IL-10*) play a role in an organism's inflammatory processes. *Mx*, the Interferon (IFN)-inducible gene, acts as a first line of defense against viral infection by blocking early steps in the viral replication cycle. In a previous study, large yellow croakers were fed an optimal dose of sitosterol, which resulted in reduced *IL-1* β levels.¹¹ Previous studies reported that Pseudomonas aeruginosa lung infection and inflammation in mice were alleviated by β sitosterol.¹⁰ Moreover, β -sitosterol significantly increased the production of antimicrobial peptides in intestinal epithelial cells and reduced the survival of intracellular Salmonella typhimurium in mice.⁹ In the present study, the expression levels of *IL-10*, *IL-1β*, *TNF-α*, *IRF-3*, and *Mx* presented different fluctuation levels in β -sitosterol treatment groups. Surprisingly, the β -sitosterol supplementation in feed had a relatively lower protection rate against *A. hy-drophila* (40%) or LMBRaV (20%) infection in largemouth bass. This may be attributed to the immune-related gene expression, or that the effects of β -sitosterol are not as prominent in disease resistance as they are in regulating growth in largemouth bass. Nevertheless, these findings require further research for confirmation.

In conclusion, in this study, dietary supplementation of 20-160 mg/kg of β -sitosterol improved growth performance, disease resistance, and antioxidant enzyme activity in largemouth bass, revealing an optimal dose of 40 mg/kg. β -sitosterol is an effective plant extract that promotes healthy fish growth.

ACKNOWLEDGMENTS

This research was funded by the National Key Research Development Program of China (2023YFD2400704), the Technical Innovation Special Project of Hubei Province (2022BBA0054, 2023BBB122), the Central Public-interest Scientific Institution Basal Research Fund (YFI 202207, 2023TD46), the National Natural Science Foundation of China (32202980) and National Freshwater Aquatic Germplasm Resource Center (FGRC18537).

AUTHORS' CONTRIBUTION

Writing – original draft: Yangyang Xing (Equal), Liping Zhang (Equal). Software: Yangyang Xing (Lead). Validation: Yangyang Xing. Project administration: Liping Zhang (Lead). Visualization: Liping Zhang (Lead). Conceptualization: Mingyang Xue (Lead). Investigation: Mingyang Xue (Lead). Methodology: Wei Liu (Lead). Formal Analysis: Nan Jiang (Lead). Funding acquisition: Yiqun Li (Lead). Data curation: Jianwu Chen (Lead). Writing – review & editing: Yan Meng (Lead).

COMPETING OF INTEREST - COPE

The authors declare that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper.

ETHICAL CONDUCT APPROVAL - IACUC

All the experimental procedures were approved by the Yangtze River Fisheries Research Institute's Animal Experimental Ethical Committee (YFI 2022-Mengyan-0412).

INFORMED CONSENT STATEMENT

All authors and institutions have confirmed this manuscript for publication.

DATA AVAILABILITY STATEMENT

The data that support the findings of this study are available from the corresponding author upon reasonable request.

Submitted: March 14, 2024 CDT, Accepted: April 22, 2024 CDT

This is an open-access article distributed under the terms of the Creative Commons Attribution 4.0 International License (CCBY-NC-SA-4.0). View this license's legal deed at https://creativecommons.org/licenses/by-nc-sa/4.0 and legal code at https://creativecommons.org/licenses/by-nc-sa/4.0/legalcode for more information.

REFERENCES

1. Huang XL, Zhong L, Kang Q, et al. A high starch diet alters the composition of the intestinal microbiota of largemouth bass micropterus salmoides, which may be associated with the development of enteritis. *Front Microbiol.* 2021;12:6965588. doi:10.3389/fmicb.2021.696588

2. Hossain MS, Small BC, Kumar V, Hardy R. Utilization of functional feed additives to produce cost-effective, ecofriendly aquafeeds high in plantbased ingredients. *Rev Aquacult*. Published online 2023:1-33. doi:10.1111/raq.12824

3. Luo XW, Shen JY, Yang T, et al. Isolation and identification of largemouth bass ranavirus from Hubei Province. *Journal of Fishery Sciences of China*. 2022;29(3):494-502.

4. Jiang N, Shen JY, Zhou Y, et al. Development of a droplet digital PCR method for the sensitive detection and quantification of largemouth bass ranavirus. *J F Dis.* 2023;46(2):91-98. <u>doi:10.1111/jfd.13721</u>

5. Ng WK, Koh CB. The utilization and mode of action of organic acids in the feeds of cultured aquatic animals. *Rev Aquacult*. 2017;9(4):342-368. doi:10.1111/raq.12141

6. García Beltrán JM, Esteban MÁ. Nature-identical compounds as feed additives in aquaculture. *Fish Shellfish Immunol*. 2022;123:409-416. doi:10.1016/j.fsi.2022.03.010

7. Beltrán JMG, Esteban MA. Nature-identical compounds as feed additives in aquaculture. *Fish Shellfish Immunol*. 2022;123:409-416. <u>doi:10.1016/j.fsi.2022.03.010</u>

8. Loizou S, Lekakis I, Chrousos GP, Moutsatsou P. Beta-sitosterol exhibits anti-inflammatory activity in human aortic endothelial cells. *Mol Nutr Food Res*. 2010;54(4):551-558. <u>doi:10.1002/mnfr.200900012</u>

9. Ding K, Tan YY, Ding Y, et al. β -Sitosterol improves experimental colitis in mice with a target against pathogenic bacteria. *J Cell Biochem*. 2018;120(4):5687-5694. doi:10.1002/jcb.27853

10. Rossi A, Bragonzi A, Medede M, et al. β-sitosterol ameliorates inflammation and Pseudomonas aeruginosa lung infection in a mouse model. *J Cyst Fibros*. 2023;22(1):156-160. <u>doi:10.1016/</u> j.jcf.2022.08.005 11. Song Y, Lu Z, Zhang X, et al. Dietary β -sitosterol supplementation enhanced intestinal immune function of large yellow croaker (Larimichthys crocea) infected with Aeromonas hydrophila. *Aquac Res.* 2022;53(18):6545-6561. doi:10.1111/are.16123

12. Zhang P, Liu NC, Xue MY, et al. Anti-Inflammatory and antioxidant properties of β -Sitosterol in copper sulfate-induced inflammation in zebrafish (Danio rerio). *Antioxidants*. 2023;12(2):391. doi:10.3390/antiox12020391

13. Gong H, Sun FG, Huang ZY, Pan ZC, Li ZH, Zou SL. Effects of phytosterol on growth performance, physical indicators,muscle composition and hepatic biochemical index of Micropterus salmoides. *Journal of Aquaculture*. 2020;41(8):26-30. <u>doi:10.3969/</u>j.issn.1004-2091.2020.08.006

14. Pan ZC, Li ZH, Sun FG, et al. Effect of phytosterols on growth performance, digestive enzymes, muscle quality and amino acid composition of Rhododendron brevicornis (Litopenaeus vannamei). *Jiangxi Fishery Science and Technology*. 2019;0(6):3-7. <u>doi:10.3969/</u> j.issn.1006-3188.2019.06.002

15. Shi Y, Cao XL, Zhong L, et al. Application of sunflower meal in diets ofd on-growing grass carp (Ctenopharyngodon idellus) and evaluation of enzymatic hydrolysis. *Aquaculture*. 2023;563:738908. doi:10.1016/j.aquaculture.2022.738908

16. Yi WY, Zhang X, Zeng K, et al. Construction of a DNA vaccine and its protective effect on largemouth bass (Micropterus salmoides) challenged with largemouth bass virus (LMBV). *Fish Shellfish Immunol*. 2020;106:103-109. <u>doi:10.1016/j.fsi.2020.06.062</u>

17. Nie YF, Wu J, Ma YP, et al. Molecular cloning of tumor necrosis factor-alpha induced protein 2 gene and expression analysis of immune-related genes associated with it in largemouth Bass (Micropterus salmoides). *Journal of Guangdong Ocean University*. 2022;42(3):1-10. doi:10.3969/ j.issn.1673-9159.2022.03.001

18. Naji TAA, Amadou I, Zhao RY, Tang X, Shi YH, Le GW. Effects of phytosterol in feed on growth and related gene expression in muscles of broiler chickens. *Trop J Pharm Res.* 2014;13(1):9. doi:10.4314/tjpr.v13i1.2

19. Hu QL, Zhuo Z, Fang SL, Zhang YW, Feng J. Phytosterols improve immunity and exert antiinflammatory activity in weaned piglets. *J Sci Food Agr*. 2017;97(12):4103-4109. doi:10.1002/jsfa.8277 20. Li ZH, Huang ZY, Pan ZC, Sun FG, Zhao HH. Effects of phytosterol on growth performance, serum lipid metabolism indicators and hepatopancreas antioxidant indicators of tilapia (Oreochromis niloticus). *Chinese Journal of Animal Nutrition*. 2019;31(12):5866-5872. <u>doi:10.3969/</u> j.issn.1006-267x.2019.12.052

21. Zhang YW, Ruan JJ, Wen C, Liu WB, Zhou YM. Effects of phytosterol on growth performance, plasma biochemical parameters and antioxidant parameters in hepatopancreas of bluntnose black bream megalobrama amblycephala. *Fisheries Science*. 2012;31(4):192-196. doi:10.16378/ j.cnki.1003-1111.2012.04.004 22. Panda S, Kar A, Patil S. Soy sterols in the regulation of thyroid functions, glucose homeostasis and hepatic lipid peroxidation in mice. *Food Res Int.* 2009;42(8):1087-1092. <u>doi:10.1016/</u>j.foodres.2009.05.001

23. Yuan CC, Ding XQ, Jiang L, Ye WX, Xu JH, Qian LC. Effects of dietary phytosterols supplementation on serum parameters, nutrient digestibility and digestive enzyme of white feather broilers. *Ital J Anim Sci.* 2021;20(1):2102-2109. <u>doi:10.1080/</u>1828051x.2021.2000895

24. Liu HQ, Guo SH, Shen YJ, Li XM. Study in antioxidative effect of β -sitosteroll. *Academic Journal of Guangdong College of Pharmacy*. 2004;20(3):281-283. <u>doi:10.16809/</u> j.cnki.1006-8783.2004.03.033