

# The Open Access Israeli Journal of Aquaculture – Bamidgeh

As from **January 2010** The Israeli Journal of Aquaculture - Bamidgeh (IJA) will be published exclusively as **an on-line Open Access (OA)** quarterly accessible by all AquacultureHub (<http://www.aquaculturehub.org>) members and registered individuals and institutions. Please visit our website (<http://siamb.org.il>) for free registration form, further information and instructions.

This transformation from a subscription printed version to an on-line OA journal, aims at supporting the concept that scientific peer-reviewed publications should be made available to all, including those with limited resources. The OA IJA does not enforce author or subscription fees and will endeavor to obtain alternative sources of income to support this policy for as long as possible.

## Editor-in-Chief

Dan Mires

## Editorial Board

<b>Rina Chakrabarti</b>	Aqua Research Lab, Dept. of Zoology, University of Delhi, India
<b>Angelo Colorni</b>	National Center for Mariculture, IOLR Eilat, Israel
<b>Daniel Golani</b>	The Hebrew University of Jerusalem Jerusalem, Israel
<b>Hillel Gordin</b>	Kibbutz Yotveta, Arava, Israel
<b>Sheenan Harpaz</b>	Agricultural Research Organization Beit Dagan,
<b>Gideon Hulata</b>	Agricultural Research Organization Beit Dagan,
<b>George Wm. Kissil</b>	National Center for Mariculture, IOLR, Eilat, Israel
<b>Ingrid Lupatsch</b>	Swansea University, Singleton Park, Swansea, UK
<b>Spencer Malecha</b>	Dept. of Human Nutrition, Food & Animal Sciences, CTAHR, University of Hawaii
<b>Constantinos Mylonas</b>	Hellenic Center for Marine Research, Crete, Greece
<b>Amos Tandler</b>	National Center for Mariculture, IOLR Eilat, Israel
<b>Emilio Tibaldi</b>	Udine University Udine, Italy
<b>Jaap van Rijn</b>	Faculty of Agriculture, The Hebrew University of Jerusalem, Israel
<b>Zvi Yaron</b>	Dept. of Zoology, Tel Aviv University, Tel Aviv, Israel

Published under auspices of  
**The Society of Israeli Aquaculture and  
Marine Biotechnology (SIAMB),  
University of Hawai'i at Mānoa Library**

&

**University of Hawai'i at Mānoa  
Aquaculture Program**  
in association with  
**AquacultureHub**

<http://www.aquaculturehub.org>



UNIVERSITY  
of HAWAII  
MĀNOA  
LIBRARY



**AquacultureHub**  
educate • learn • share • engage

ISSN 0792 - 156X

© Israeli Journal of Aquaculture - BAMIGDEH.

PUBLISHER:

Israeli Journal of Aquaculture - BAMIGDEH -  
Kibbutz Ein Hamifratz, Mobile Post 25210,  
ISRAEL


Phone: + 972 52 3965809

<http://siamb.org.il>

Copy Editor **Ellen Rosenberg**



The IJA appears exclusively as a peer-reviewed on-line open-access journal at <http://www.siamb.org.il>. To read papers free of charge, please register online at [registration form](#).  
Sale of IJA papers is strictly forbidden.



## Oxygen Consumption and Lethal Dissolved Oxygen Level of Hybrids of *Siniperca chuatsi*♀ × *S. scherzeri*♂

Jijia Sun<sup>1,2</sup>, Jianguo He<sup>1</sup>, Guifeng Li<sup>1\*</sup>, Li Liu<sup>2</sup>, Xiaocui Hou<sup>1</sup>, Pengfei Wang<sup>1</sup>

<sup>1</sup> School of Life Sciences, Sun Yat-sen University, Guangzhou 510275, P.R. China

<sup>2</sup> College of Animal Science, South China Agriculture University, Guangzhou 510642, P.R. China

(Received 19.3.2015, Accepted 21.6.2015)

Key words: hybrid, *Siniperca chuatsi*, *S. scherzeri*, oxygen consumption rate, lethal dissolved oxygen level, circadian rhythm

### Abstract

Oxygen consumption rate (OCR) and lethal dissolved oxygen levels (DO) of the hybrids (225.3 ± 4.6 g) of *Siniperca chuatsi* ♀ × *S. scherzeri* ♂, were determined after acclimating the fish to 15, 20, 26 and 30°C for 20 days. The same parameters were measured for hybrids with different body weight (146.8 ± 6.3 g, 234.3 ± 8.4 g, 273.9 ± 3.3 g, 327 ± 5.1 g) at 26 °C. OCR increased significantly ( $P < 0.05$ ) with increasing acclimation temperature between 20-26°C, and decreased with increasing body weight. Final preferred temperature estimated from the  $Q_{10}$  was between 26-30°C. The lethal DO concentration for the hybrids at 30°C was significantly higher ( $P < 0.05$ ) than at other temperatures. OCR over the daily cycle under natural lighting conditions was also determined at 26°C. It showed a circadian rhythm with the lowest point at 0830 and peak rates at 1430. Results show that the hybrid has a higher stress tolerance and higher stocking density than the female parent *S. chuatsi* and indicate their culture potential in subtropical freshwater regions.

\* Corresponding author. E-mail: [liguif@mail.sysu.edu.cn](mailto:liguif@mail.sysu.edu.cn)

## Introduction

The measurement of oxygen consumption has been proved to be an efficient method of estimating the metabolic rate of fish, and several different states of respiratory metabolism have been classified for fish as standard, routine, active, and anerobic metabolic rate (Brett et al., 1979; Cech, 1990). Oxygen consumption rate (OCR) of aquatic organisms is one of the physiological responses that correlate with changes in environmental factors, such as temperature, light intensity, dissolved oxygen, and salinity (Salvato et al., 2001). Energy expenditure associated with physiological stress that these factors impose on organisms can also be estimated by measuring OCR (Altinok et al., 2003; Brougher et al., 2005).

*Siniperca chuatsi* (Basilewsky) and *Siniperca scherzeri* (Steindachner) are the most economically and geographically important siniperids for aquaculture in China (Zhou et al., 1988). In recent years, studies have shown that the hybrids of *S. chuatsi* ♀ × *S. scherzeri* ♂ combined the rapid growth trait of the female parent and the body surface coloration of the male parent (Wang et al., 2013). Morphological analysis (Mi et al., 2010) and DNA marker (SSR and SNP) studies have been carried out on this hybrid (Wang et al., 2013; Qu et al., 2013; Huang et al., 2013). However, the effects of hybridization on oxygen consumption of this hybrid have not been reported.

In this laboratory study, the effects of temperature and body weight on oxygen consumption of the hybrid were investigated, and the circadian rhythm of OCR and the lethal (DO) levels were measured.

## Materials and Methods

**Fish.** The hybrids of *Siniperca chuatsi* ♀ × *S. scherzeri* ♂ were obtained from Shengsheng Fisheries Ltd. at Foshan in Guangdong Province, China. In the laboratory, 60 fish with wet body weight ranging from 150 to 350 g were placed in a reservoir with constant aeration. Hybrids were starved for 24h before commencing the experiment to reduce associated metabolic responses (Nelson et al., 1985; Beamish et al., 1990).

**Experimental facilities.** Oxygen consumption rate of the hybrids was determined using a flow-through respirometer (Fig. 1). Hybrids were kept in the respiratory chamber (250×250×300 mm), which was immersed in a 300 L water bath (1500×550×350 mm) for temperature control. The constant temperature treatments of the respirometer were achieved by a mixed temperature control system, which included a thermostat controlling the on/off switch of a 1000 W electric heater, and a manual controller.

**Determination of oxygen consumption rates.** Two OCR measurement experiments were implemented. (1) OCR of hybrids was measured at four acclimation temperatures (15, 20, 26 and 30°C). The mean wet body weight of the fish was  $225.3 \pm 4.6$  g and there were no significant differences in wet weights among the treatments ( $P > 0.05$ ). Fish were maintained at predetermined temperatures for 20 days to ensure full thermal adaptation. Prior to the determination of oxygen consumption, water flow was controlled at 6.0 L/h. The oxygen content of outflow water was  $\geq 5$  mg/L. All four sides of the chamber were covered with opaque screens to minimize photoperiodic effects on the experimental fishes (fish?). When fish became quiescent after 2 h, oxygen consumption was determined over a period of 6 h and water in the chamber was siphoned off every 2h. Eight fish from each acclimation temperature were introduced one at a time into the chamber. For each treatment, there was a blank control to correct for the respiration of




Fig. 1. Diagram of a flow-through respirometer: 1.water tank; 2.water level controller; 3.siphon tube; 4.thermometer; 5.rubber plug; 6.respiratory chamber; 7.water bath; 8.control valve; 9.Sampling vessel.

bacteria in the water. Measurements were made between 0800 and 1200 hours. Oxygen content of water samples was determined using the Winkler method (Strickland et al., 1968). The OCR was calculated using the following formula:

$$\text{OCR (mg O}_2\text{ / kg/h)} = (\text{DO}_1 - \text{DO}_2) \times V/W$$

- (1) where  $\text{DO}_1$  is the concentration of oxygen ( $\text{mg O}_2/\text{L}$ ) in the chamber without fish,  $\text{DO}_2$  is the concentration of oxygen in chamber with fish,  $V$  is the water flow rate ( $\text{L/h}$ ),  $W$  is the wet body weight of hybrid ( $\text{Kg}$ ).

The temperature coefficients ( $Q_{10}$ ) were calculated to assess the effect of temperature acclimation on OCR by using the formula (Schmidt-Nielsen, 1997):

$$Q_{10} = (\text{Rate}_2 / \text{Rate}_1)^{10/(\text{Temp}_2 - \text{Temp}_1)}$$

- (2) OCRs of four different body weight groups ( $146.8 \pm 6.3$  g,  $234.3 \pm 8.4$  g,  $273.9 \pm 3.3$  g, and  $327 \pm 5.1$  g) were measured at  $26^\circ\text{C}$ . Six fish from each weight group were individually introduced into the chamber. The procedure followed was then identical to that of the acclimation temperatures.

*Measurement of lethal dissolved oxygen (DO) concentration.* After the oxygen consumption measurements, the water flow was shut off and the test fish were kept in the sealed respiratory chamber. To determine the lethal DO level, the DO concentration in the chamber was measured for a final time just after the death of each fish. After each lethal DO level was measured, the body weight of dead fish was measured immediately.

*Circadian rhythm in the oxygen consumption.* After removal of the opaque screens, the chamber was exposed to a natural photoperiod. In this test, the mean temperature was  $26 \pm 1^\circ\text{C}$  and the body weight of the fish used was 136.8 g, 168.2 g, 203.7 g, and 285.4 g, respectively. Measurements of OCR were made, every 2 h during the 24 period. light period from 0630 to 1630 hours, and dark period was from 1830 to 0430 hours.

*Statistical analysis.* The data were analyzed by SPSS for Windows (Version 17.0) and expressed as mean  $\pm$  SE. Inter-treatment differences for OCR and lethal DO concentration were analyzed with one-way ANOVA followed by least significant difference (LSD) multiple range tests. Differences were considered statistically significant when  $P < 0.05$ .

## Results

The OCR of the hybrids varied with the acclimation temperature (Fig. 2). The OCR changed significantly ( $P < 0.05$ ) as the acclimation temperature increased from  $15^\circ\text{C}$ - $26^\circ\text{C}$ , first decreasing to a minimum rate of  $92 \text{ mg O}_2/\text{Kg/h}$  at  $20^\circ\text{C}$  and then reaching a peak rate of  $136.8 \text{ mg O}_2/\text{Kg/h}$  at  $26^\circ\text{C}$ . However, no significant change ( $P > 0.05$ ) in the OCR was observed between  $26^\circ\text{C}$ - $30^\circ\text{C}$ . Maximum and minimum temperature quotients  $Q_{10}$  were observed between  $20^\circ\text{C}$ - $26^\circ\text{C}$  and  $15^\circ\text{C}$ - $20^\circ\text{C}$  as 1.94 and 0.65, respectively. There was a drop in the  $Q_{10}$  when the acclimation temperature increased from  $26^\circ\text{C}$ - $30^\circ\text{C}$ . The OCR decreased with increasing body weight (Fig. 3), and the OCR at mean body weight of  $327.0 \pm 5.1$  g was significantly lower than the others ( $P < 0.05$ ).

The effects of acclimation temperature and body weight on the lethal DO concentration are showed in Fig. 4. Among the four experimental temperature treatments, lethal DO levels increased with increasing temperature. The lethal DO concentration reached a maximum of  $1.0648 \pm 0.1244 \text{ mg/L}$  at  $30^\circ\text{C}$ , and this value was significantly higher than at other temperatures ( $P < 0.05$ ) [Fig. 4(a)]. Lethal DO levels decreased in relation to body weight increase, except for a significant increase from  $273.9 \pm 3.3$  g to  $327.0 \pm 5.1$  g ( $P < 0.05$ ) [Fig. 4(b)].

The circadian rhythms in the OCRs of the four different body weight hybrid groups are similar (Fig. 5). The lowest point in OCR was recorded at 0830 hours (the mean OCR  $32.1 \text{ mg O}_2/\text{Kg/h}$ ), and the peak rate in OCR at 1430 hours (the mean OCR  $178.1 \text{ mg O}_2/\text{Kg/h}$ ). The mean OCRs during the day (148.0, 132.6, 147.3 and  $109.9 \text{ mg O}_2/\text{Kg/h}$  for body weight 136.8, 168.2, 203.7 and 285.4 g respectively) were lower than at night (181.9, 141.1, 163.5 and  $125.0 \text{ mg O}_2/\text{Kg/h}$ ), which show a significant circadian rhythm.




Fig. 2. OCR of hybrids acclimated to 15, 20, 26 and 30°C. Values are expressed as mean  $\pm$  SE. Different letters on the value indicate significant difference ( $P < 0.05$ ). Numbers in parenthesis are  $Q_{10}$  between acclimation temperatures (15-20, 20-26 and 26-30°C).




Fig. 3. OCR of hybrids with different body weight. Values are expressed as mean  $\pm$  SE. Different letters on the value indicate significant difference ( $P < 0.05$ ).




Fig. 4. (a) Lethal DO concentration at different temperature; (b) Lethal DO concentration at different body weight. Different letters over the value indicate significant difference ( $P < 0.05$ ).




Fig. 5. The circadian rhythm in the OCRs of the four different body weight hybrids

### Discussion

In poikilothermic animals, oxygen consumption is often used as an indirect measure of metabolism and is strongly dependent on acclimation temperature (Kita et al., 1996). For many fish, such as *Lepomis macrochirus* and *L. megalotis* (Dent et al., 2003), advanced fingerlings of Indian Major Carps (*Labeo rohita*, *Catla catla* and *Cirrhinus mrigala*) (Das et al., 2004), *Pangasius pangasius* (Debnath et al., 2006), and *Micropterus salmoides* juveniles (Díaz et al., 2007), a linear correlation has been reported between oxygen consumption and acclimation temperature. However, some researchers have suggested that metabolic rates of some fish are negatively related to acclimation temperature (Brown et al., 1989; Walsh et al., 1997).

In the present study, OCR of hybrids showed positive temperature dependence of metabolism from 20°C-26°C and negative temperature dependence from 15°C-20°C and 26°C-30°C. Compared with the OCRs of *Siniperca chuatsi* (average body weight  $230 \pm 11.79$  g) at 20°C (139.0 mgO<sub>2</sub>/Kg/h), 26°C (173.7 mgO<sub>2</sub>/Kg/h) and 30°C (174.1 mgO<sub>2</sub>/Kg/h) (Si et al., 1995), the hybrids had a lower OCR level, and the mean OCR decrease was 41.4 mg O<sub>2</sub>/Kg/h. This indicates that the hybrids have a stronger stress tolerance than their parents and could be reared at higher stocking densities than *S. chuatsi*. However, the OCR of the hybrids at 15°C was significantly higher than at 20°C and higher than that of *S. chuatsi*. It seems likely that metabolism of hybrids may be more efficient at lower temperatures than *S. chuatsi*.

A freshwater teleost demonstrates physiological plasticity when it is able to regain or approach its metabolic set-point (e.g. quantitative or qualitative changes in enzyme expression) within the context of thermally fluctuating environments (Dent et al., 2003). This relationship predicts the point where the Q<sub>10</sub> for oxygen consumption starts to decrease with increasing acclimation temperatures that corresponds to their optimal temperature for growth (Kita et al., 1996). In our study, the optimal temperature for hybrids ranged from 26°C-30°C.

Our study found the OCRs of hybrids decreased with increasing body weight. This negative correlation of OCR and weight was also seen in studies on *S. chuatsi* (Si et al., 1995). Furthermore, the lethal DO levels and the circadian rhythm of the hybrids were similar to those of *S. chuatsi* (Si et al., 1995). These results demonstrated that hybrids had similar respiratory physiological characteristics to their female parent. However, our study also demonstrated that the hybrids had some distinctive oxygen consumption characteristics. The lethal DO concentration for hybrids at 30°C was significantly higher than at other temperatures and nearly 1.7 times higher than that of *S. chuatsi* (Si et al., 1995). The lowest levels in OCR of hybrids occurred at 0830 and 1430, while lowest levels for *S. chuatsi* (female parent) and *S. scherzeri* (male parent) occurred at 0600 and 1200 and 0700 and 1200 (Si et al., 1995; Peng et al., 2005), respectively.

In conclusion, the present study showed that the hybrids had higher stress tolerance and higher stocking density than their female parent *S. chuatsi* from 20°C-30°C. In addition, the final preferred temperature (26°C-30°C) of hybrids based on the Q<sub>10</sub> value indicates higher culture potential in subtropical freshwater regions. Future researches on various biochemical changes of hybridization, such as metabolic enzymes and heat shock protein, may aid in the understanding of differences in metabolic mechanisms of the hybrid.

### Acknowledgements

This research was supported by a project of the National Key Technology R&D Program, China (No. 2012BAD25B04), a project of the Science and Technology Planning Project of Guangdong Province, China (No. 2012A020800001), a project of the Educational Commission of Guangdong Province, China (No. cxzd1104), and a Cooperative Project of Guangdong Province, China (No. 2011B090400179).

### References

- Altinok I. and J.M. Grizzle**, 2003. Effects of low salinities on oxygen consumption of selected euryhaline and stenohaline freshwater fish. *J World Aquacult Soc*, 34(1): 113-117.
- Beamish F.W.H. and E.A. Trippel**, 1990. Heat Increment: a static or dynamic dimension in bioenergetic models? *Transactions of the American Fisheries Society*, 119(4): 649-661.
- Brett J.R. and T.D.D Groves**, 1979. Physiological energetics. pp. 279-352. In: W.S. Hoar, D.J. Randall, and J.R. Brett (eds.). *Fish Physiology, Vol VIII: Bioenergetics and Growth*. Academic Press, New York, USA.
- Brougher D.S., Douglass L.W. and J.H. Soares**, 2005. Comparative oxygen consumption and metabolism of striped bass *Morone saxatilis* and its hybrid *M. chrysops* ♀ × *M. saxatilis* ♂. *J World Aquacult Soc*, 36(4): 521-529.
- Brown L.R.**, 1989. Temperature preferences and oxygen consumption of three species of sculpin (Cottus) from the Pit River drainage, California. *Environ Biol Fish*, 26: 223-236.
- Cech J.J.**, 1990. Respirometry. pp. 335-362. In: C.B. Schreck, P.B. Moyle (eds.). *Methods Fish Biol*. American Fisheries Society, Bethesda, MD, USA.
- Das T., Pal, A.K., Chakraborty S.K., Manush S.M., Chatterjee N. and S.C. Mukherjee**, 2004. Thermal tolerance and oxygen consumption of Indian Major Carps acclimated to four temperatures. *J Thermal Biol*, 29(3): 157-163.
- Debnath D., Pal A.K., Sahu N.P., Baruah K., Yengkokpam S., Das T. and S. Manush**, 2006. Thermal tolerance and metabolic activity of yellowtail catfish *Pangasius pangasius* (Hamilton) advanced fingerlings with emphasis on their culture potential. *Aquaculture*, 258(4): 606-610.
- Dent L. and W.I. Lutterschmidt**, 2003. Comparative thermal physiology of two sympatric sunfishes (Centrarchidae: Perciformes) with a discussion of microhabitat utilization. *J Thermal Biol*, 28(1): 67-74.
- Díaz F., Denisse Re A., González R.A., Sánchez L.N., Leyva G. and F. Valenzuela**, 2007. Temperature preference and oxygen consumption of the largemouth bass *Micropterus salmoides* (Lacépède) acclimated to different temperatures. *Aquacult Res*, 38(13): 1387-1394.
- Huang W., Liang X. and Qu C.M.**, 2013. Development and characterization of novel polymorphic microsatellite loci in *Siniperca scherzeri* Steindachner and *Siniperca chuatsi* (Basilewsky). *Molec Biol Reports*, 40(2): 751-756.
- Kita J., Tsuchida S. and T. Setoguma**, 1996. Temperature preference and tolerance, and oxygen consumption of the marbled rock-fish, *Sebasticus marmoratus*. *Mar Biol*, 125: 467-471.
- Mi G., Zhao J., Jia Y., Deng Y., Lian Q. and Y. Wang**, 2010. Morphological and microsatellite analysis of *Siniperca chuatsi* ♀ × *Siniperca scherzeri* ♂ hybrid with their parents. *Journal of Shanghai Ocean University*, 19(2): 145-150 (in Chinese with English abstract).
- Nelson S.G., Simmons M.A. and A.W. Knight**, 1985. Calorigenic effect on diet on the grass shrimp *Crangon franciscorum* (Crustacea: Crangonidae). *Comp Biochem Physiol A: Physiol*, 82(2): 373-376.
- Peng Z., Zeng K., Gao Y., Zhu S. and H. Zhang**, 2005. Studies on the oxygen consumption rate and suffocation point of *Siniperca Scherzeri* fingerling. *Inland Aquatic Product*, 30(4): 30-33 (in Chinese).
- Qu C.M., Liang X., Huang W., Zhao C., Liang C., Yang M. and C. Tian**, 2013. Development and characterization of twenty-nine novel polymorphic microsatellite loci in the mandarin fish *Siniperca chuatsi*. *J Genetics*, 92(1): 19-23.
- Salvato B., Cuomo V., Di Muro R. and M. Beltramini**, 2001. Effects of environmental parameters on the oxygen consumption of four marine invertebrates: a comparative factorial study. *Marine Biology*, 138(4): 659-668.
- Schmidt-Nielsen K.**, 1997. *Animal Physiology: Adaptation and Environment*. Cambridge University Press, UK.



- Si Y., Chen Y. and J. Zeng**, 1995. The oxygen consumption and cultivation of Mandarin fish, *Siniperca Chuatsi*. *Acta Hydrobiologica Sinica*, 19(4): 327-332 (in Chinese with English abstract).
- Strickland J.D.H. and T.R. Parsons**, 1968. A practical handbook of seawater analysis. *Bulletin (Fisheries Research Board of Canada)*, 167:1-11.
- Walsh S.J., Haney D.C. and C.M. Timmerman**, 1997. Variation in thermal tolerance and routine metabolism among spring and stream-dwelling freshwater sculpins (Teleostei: Cottidae) of the southeastern United States. *Ecol Freshwater Fish*, 6(2): 84-94.
- Wang H., Sun J., Lu X., Wang P., Xu P., Zeng L., Yu D. and G. Li**, 2013. Identification of insulin-like growth factor I gene polymorphisms using high-resolution melting and its effect on growth traits in sinipercid species. *Fish Sci: FS*, 79(3):439-446.
- Zhou C.W., Yang Q. and D.L. Cai**, 1988. On the classification and distribution of the sinipercinae fishes (family Serranidae). *Zoological Res*, 9(2):113-125 (in Chinese with English abstract).