

## Original Research Articles

# Age, growth, and mortality of Bogue (*Boops boops*, Linnaeus, 1758) from the Antalya Bay (Northwest Mediterranean Sea, Türkiye)

Habil U. Koca<sup>1</sup><sup>1</sup> Department of Fishing and Processing Technology, Faculty of Eğirdir Fisheries, Isparta University of Applied Sciences, PC 32260, Isparta, Türkiye

Keywords: Bogue, population structure, Antalya Bay, Türkiye

<https://doi.org/10.46989/001c.84080>

## Israeli Journal of Aquaculture - Bamidgeh

Vol. 75, Issue 2, 2023

Bogue (*Boops boops*, Linnaeus, 1758) that was gathered in Antalya Bay had their ages, growth rates, and mortality examined. Using the von Bertalanffy equation,  $L_t = 37.50 (1 - e^{-0.15(t + 1.32)})$  ( $R^2 = 0.988$ ), growth in length was expressed for the entire sample without revealing any disparities between the sexes. The values of the allometric coefficient ( $b$ ) of the length-weight relationships specify negative allometric growth ( $b < 3$ ) in females, males, and overall sexes. Total and natural mortality were found to be  $Z = 1.12 \text{ y}^{-1}$ , and  $M = 0.36 \text{ y}^{-1}$ . The exploitation ratio  $E = 0.68$  indicated that the population was overexploited.

## INTRODUCTION

Bogue (*Boops boops*, Linnaeus, 1758), family Sparidae, is an important demersal and semi-pelagic species in the Mediterranean Sea. This species lives in the Eastern Atlantic, in an area reaching from Norway in the north, where it is occasionally to Angola in the south and throughout the Mediterranean Sea including the Black Sea.<sup>1</sup> Capture production of bogue in Turkey ranks 10th with an amount of 2310.6 tones in 2022 after anchovy, Atlantic bonito, pilchard, mackerel, sprat, whiting, blue fish, twaite shad, and chup mackerel.<sup>2</sup>

Age, growth, and mortality of *B. boops*, have been studied from the Portuguese coast, Egyptian waters, western Mediterranean coast of Egypt, Tunisian waters, Algarve (South of Portugal), northern Aegean Sea, İzmir Bay, Adriatic coasts (Velebit Channel), Egyptian Mediterranean waters off Alexandria Egypt, İzmir Bay, Montenegrin waters, northern and central Adriatic Sea, Gulf of Antalya, southern Tyrrhenian Sea, between Nador and Sidia (Morocco), Güllük Bay, Mediterranean waters front Alexandria, Algerian East coast, and Saros Bay.<sup>3-22</sup> Despite its commercial importance, the biology and population parameters of *B. boops* are not well known in the northeastern Mediterranean Sea, Türkiye.

The purpose of this study was to investigate the age, growth, and mortality of the *B. boops* species from the Gulf of Antalya. Results are important for future improvement and sustainable fishery management plans for this species.

## MATERIALS AND METHODS

The samples of *B. boops* were collected monthly from the commercial landings from October 2020 to September 2021 from the Gulf of Antalya (Figure 1). Samplings of fish were made by a bottom trawl net with a 44 mm mesh size. Total length was measured to the nearest 0.1 mm and body weight to the nearest 0.01 g. Sex was determined by macroscopic observation of the gonads. Gonads of all individuals were dissected and weighed to the nearest 0.01 g. Age determination was carried out by reading annual rings on *B. boops* right and left sagittal otoliths. The number of opaque zones and the presence of marginal translucent zone were checked by two readers.

The growth, condition factor, length-weight relationship, and exploitation were statistically analyzed using FISAT II as detailed in detail by Gayanilo et al.<sup>23</sup> Length and weight relative growth rate were calculated by following equations Ricker.<sup>24</sup> The relative growth in length ( $(L_2 - L_1 / L_1) \times 100$ ), the relative growth in weight ( $(W_2 - W_1 / W_1) \times 100$ ), where  $L_1$ : the mean length of the first age group (cm),  $L_2$ : the mean length of the next age group (cm),  $W_1$ : the mean weight of the first age group (g),  $W_2$ : the mean weight of the next age group (g).

The condition factor ( $K$ ) determined changes in the species' nutritional condition.  $K$  was estimated using the equation proposed by Ricker<sup>24</sup>:  $K = (W / L^3) \times 100$ , where  $W$  = Total weight of fish (g),  $L$  = Total length of fish (cm). The gonad-somatic index ( $GSI$ ) was calculated using the  $GSI = W_g / W \times 100$ ; where  $W_g$  is gonad weight and  $W$  is body weight. The length-weight relationship was described by the equation:  $W = aL^b$ , where  $W$  is the weight of fish (g),  $L$  is

a \* Corresponding author. e-mail: [ugurkoca@isparta.edu.tr](mailto:ugurkoca@isparta.edu.tr)




Figure 1. *B. boops* sample and map of the study area

the total length (cm),  $b$  is the allometric rate and  $a$  is a constant.<sup>25</sup> The linear regression relationship for females and males was tested for significance using analysis of variance (ANOVA).

The von Bertalanffy growth model was fitted to the observed lengths at age. The model is described by the equation  $L_t = L_\infty (1 - e^{-k(t-t_0)})$ ,<sup>24</sup> where  $L_\infty$  is the asymptotic total length,  $L_t$  is the total length at age  $t$ ,  $k$  is the growth curvature parameter and  $t_0$  is the theoretical age that fish would have been at zero length.<sup>26</sup> The growth performance index ( $\phi$ ) was computed from the expression given by Pauly and Munro<sup>27</sup>:  $\phi = \log_{10} k + 2 \log_{10} L_\infty$ .

The total mortality rate ( $Z$ ) was estimated by using the length converted catch curve. The natural mortality rate ( $M$ ) was estimated using by the equation of Pauly<sup>28</sup>:  $\log M = -0.0066 - 0.279 \log L_\infty + 0.6543 \log k + 0.4634 \log T$ . Where  $T$  is the mean annual temperature, estimated in the study area at 22.2 °C,  $k$  and  $L_\infty$  are the constants of the von Bertalanffy equation. The fishing mortality rate  $F$  was estimated using the relationship of Pauly<sup>29</sup>:  $F = Z - M$ . The exploitation rate  $E$  was obtained by the relationship of Gulland<sup>30</sup>:  $E = F / Z$ . The stock is in equilibrium when  $E = 0.5$ , it is under-exploited when  $E < 0.5$  and is overexploited when  $E > 0.5$ .

A t-test was used for possible variations in the length-weight of females, and males, and the overall sex ratio was assessed using the chi-square test.<sup>31</sup> An analysis of covariance was used to compare relative growth, condition factor, and the linear regression relationship for females and males was tested for significance using analysis of variance (ANOVA).

## RESULTS

A total of 641 specimens of *B. boops* ranging from 10.8 to 24.9 cm in total length and 12.99 to 141.11 g weight were sampled. There were 316 females (10.8 - 24.9 cm, TL; 12.99 - 141.11 g, W) and 325 males (10.9 - 23.0 cm, TL; 13.47 - 119.60 g, W). Females dominated length sizes were 14.0 to 17.9 cm ( $16.13 \pm 0.06$ ), whereas males were 14.0 to 18.9 cm ( $16.09 \pm 0.07$ ) (Figure 2).

The female: male ratio was 1:0.97, it was not significantly different from the expected 1:1 ratio ( $\chi^2 = 0.126$ ,  $P > 0.05$ ). The ages of individuals ranged from 1 to 5 years. In our study, the RG rates calculated according to ages: in weight and length, for 1 to 2, 2 to 3, 3 to 4 and 4 to 5 ages: 159.25 g / 28.77 cm, 58.19 g / 21.81 cm, 29.17 g / 17.56 cm, and 18.97 g / 8.78 cm, respectively.

Figure 3 show monthly variations in the relative growth (RG) and condition factor (K) calculated from 316 females and 325 males of *B. boops*. The highest relative growth values were calculated in April (13.48) after which a decrease in relative growth values was calculated with increase of age reaching in September (1.29). Values of the condition factor are relatively low for both sexes. The highest values were noticed in April 1.55 for females and 1.69 for males and the lowest values in October 0.57 for females and 0.69 for males. No significant correlation was found between relative growth and condition factor ( $P > 0.05$ , t-test).

In the *B. boops* from the Antalya Bay, the mean monthly gonadosomatic index (GSI) values were the highest in March (2.19-2.11). Spawning occurred both females and males between January (0.98-0.93) and ends in September (0.21-0.19) (Figure 4).

The total length-weight relationships were calculated respectively for females, males and overall,  $W = 0.013$




Figure 2. Length frequency distribution of female and male *B. boops* sampled




Figure 3. Monthly variations relative growth (RG) and condition factor (K) *B. boops* from the Antalya Bay.

$TL^{2.897}$ ,  $W = 0.014 TL^{2.867}$  and  $W = 0.013 TL^{2.882}$ , respectively (Figure 5).

The slopes of the total length-weight relationships, which do not differ significantly between sexes (one-way ANOVA,  $P < 0.001$ ), indicate the negative allometric growth ( $b$  ranged 2.867 and 2.897), respectively had a significant difference from the value 3.0 ( $t$ -test,  $P > 0.05$ ).

Right and left sagittal otoliths were collected from 641 individuals of *B. boops*. The results of reading otolith rings

in females, males, and overall, of bogue are shown in Table 1. The observed lengths of individuals determined for each group were used to appropriate the von Bertalanffy growth model. The von Bertalanffy growth formulas:

in females,  $L_t = 37.14 (1 - e^{-0.15(t+1.33)})$  ( $R^2 = 0.918$ );  $W_t = 237.38 (1 - e^{-0.14(t-0.43)})$  ( $R^2 = 0.996$ ), in males,  $L_t = 29.95 (1 - e^{-0.22(t+0.95)})$  ( $R^2 = 0.881$ );  $W_t = 173.74 (1 - e^{-0.21(t-0.62)})$  ( $R^2 = 0.979$ ), and overall,  $L_t = 37.50 (1 - e^{-0.15(t+1.32)})$  ( $R^2 = 0.988$ );  $W_t = 199.56 (1 - e^{-0.17(t-0.53)})$  ( $R^2 = 0.989$ ).




Figure 4. The gonadosomatic index (GSI) values of the *B. boops*.




Figure 5. The total length-weight relationship of *B. boops* (overall).


The performance index parameters estimated for *B. boops* using the length frequency data revealed the best fit for  $L_{\infty} = 37.14$  cm,  $k = 0.15$  (in females),  $L_{\infty} = 29.95$  cm,  $k = 0.22$  (in males),  $L_{\infty} = 37.50$  cm,  $k = 0.15$  (total) per year while the growth performance was estimated as 2.32, 2.29 and 2.33, respectively.

The total mortality ( $Z$ ) of bogue estimated by the length converted catch curve was 1.12 (Figure 6). The natural mortality ( $M$ ) as per Pauly's empirical formula was found to

be 0.36. The estimated fishing mortality ( $F$ ) is 0.76 and the exploitation ratio ( $E$ ) is 0.68. The exploitation ratio  $E > 0.50$ . The values of  $E$  obtained indicated that the fishing pressure exerted on the *B. boops* in the area under study was excessive fishing (TÜİK<sup>2</sup> reported that it is around 2300 tones in 2022).

**Table 1. The results according to sexes, age, total length (cm), and weight (g) (N, min., max., mean, and standard error).**

SEX	AGE				
	1	2	3	4	5
FEMALES	23	117	145	16	15
	(10.8-13.4)	(13.7-16.2)	(16.2-21.1)	(21.2-22.5)	(22.5-24.9)
	11.92±0.158	15.24±0.061	17.55±0.105	21.83±0.109	23.29±0.220
	(12.99-24.41)	(18.57-47.83)	(33.54-87.68)	(79.25-106.96)	(108.71-141.11)
	17.54±0.699	46.24±0.640	69.30±0.997	93.33±2.149	110.56±2.549
MALES	19	138	131	28	9
	(10.9-13.3)	(13.5-16.2)	(16.2-21.0)	(21.2-22.5)	(22.5-23.0)
	11.89±0.173	15.16±0.059	17.51±0.106	21.87±0.079	22.71±0.075
	(13.47-25.94)	(17.53-48.41)	(33.30-86.75)	(78.15-115.22)	(103.57-119.60)
	17.67±0.926	46.95±0.574	68.87±1.009	94.64±1.582	107.16±1.690
OVERALL	42	255	276	44	24
	(10.8-13.4)	(13.5-16.2)	(16.2-21.1)	(21.2-22.5)	(22.5-24.9)
	11.80±0.115	15.20±0.042	18.51±0.073	21.76±0.063	23.67±0.150
	(12.99-25.94)	(17.53-48.41)	(33.30-87.68)	(78.15-115.22)	(103.57-141.11)
	17.58±0.560	46.59±0.430	69.91±0.700	94.18±1.260	109.05±1.820

**Figure 6. Catch curve of *B. boops* (x: points not used; o: points used).**

## DISCUSSION

The length and weight, however, varied from various localities (Table 2). The variable in bogue growth may result from a few factors including environmental conditions or genetic variations. There are in agreement with the findings of this study. Five age groups (1 - 5) were determined in both sexes. Two and three-year-old fish dominated the population (82.84 %). In the previous studies, age classes for bogue were determined as 1 - 6; 1 - 4; 1 - 5, and 1 - 5 by El-Haweet<sup>5</sup>; El-Okda<sup>11</sup>; Kara and Bayhan<sup>12</sup>, and Dahel et al.,<sup>10</sup> respectively. The different exploitation patterns and/or ecological conditions related to habitat quality and temperature could explain age class variations.<sup>32</sup>

The condition factor is an index that expresses the degree of well-being, relative robustness, and interactions between abiotic and biotic factors in fish physiological conditions.<sup>33</sup> In the study, the mean ratio of the condition factors

for females ( $0.97 \pm 0.007$ ), males ( $0.98 \pm 0.006$ ), and overall ( $0.97 \pm 0.005$ ) for *B. boops* were all ~1.0. No significant difference was found between the mean condition factors of individuals females and males for bogue ( $P = 0.36$ ,  $P > 0.05$ ).

The estimated  $b$  values of the regression for *B. boops* showed a negative allometric growth pattern ( $b < 3$ ) for this study. Allen<sup>34</sup> chose to keep the value of the  $b$  exponent constant at 3 for isometrical fish growth. The negative allometric growth were recorded from the Gulf of Antalya,<sup>16</sup> in the Morocco waters,<sup>18</sup> in the Egypt waters,<sup>20</sup> and in the Algerian waters,<sup>21</sup> while the positive allometric growth were in the Greece waters,<sup>35</sup> in the Adriatic Sea,<sup>15</sup> and in the Aegean Sea.<sup>13</sup> Variations in fish shape, and growth increase some of the environmental conditions like temperature, season, habitat and food presence.<sup>36</sup> The factors are affected by sequences factors including gonad maturity, stomach fullness, sex, diet, and health.<sup>37</sup>

Table 2 shows the von Bertalanffy growth parameters and the growth performance in the different localities for bogue species. The primary input data for several models used to manage overfished fish stocks were 37.50 cm, 0.15 yr<sup>-1</sup>, and 1.32 for the von Bertalanffy growth parameters ( $L_{\infty}$ ,  $K$ , and  $t_0$ ). The potential causes of these discrepancies could be the variable maximum lengths recorded in the captures or the diverse sample methodologies and calculating methods utilized by various authors at various times and locations. Therefore, geographical dispersion, changes in environmental circumstances, climate influences, quality of available food, and fishing effort would all be factors in the variability of growth.

Estimates of the total mortality ( $Z$ ), the natural mortality ( $M$ ), the fishing mortality ( $F$ ), and exploitation ratio ( $E$ ) for the presently reported species were 1.12 yr<sup>-1</sup>, 0.36 yr<sup>-1</sup>, 0.76 yr<sup>-1</sup> and 0.68 which is similar to those for other references:

**Table 2. The mean total length (cm), mean weight (g), the von Bertalanffy growth parameters, and the growth performance in the different localities for bogue.**

Localitie	Sex	TL	W	$L_{\infty}$ (cm)	k	$t_0$	$\phi$	Reference
Egypt	Overall	16.02	-	31.68	0.15	1.78	2.19	Allam <sup>4</sup>
Tunisian	Overall	-	-	28.67	0.20	1.41	-	Khemiri et al. <sup>6</sup>
Algarve	Overall	19.9	-	28.06	0.22	1.42	-	Monteiro et al. <sup>7</sup>
Egypt	Overall	13.58	28.66	30.11	0.15	1.51	2.14	El-Okda <sup>11</sup>
Algeria	Female	-	-	30.66	0.30	0.00	2.45	Bouaziz et al. <sup>40</sup>
	Male	-	-	26.66	0.30	0.00	2.33	
Morocco	Overall	-	-	30.00	0.41	0.30	-	Layachi et al. <sup>18</sup>
Türkiye	Female	19.14	75.62	30.79	0.24	0.90	2.36	Kara and Bayhan <sup>13</sup>
	Male	20.28	91.41	29.87	0.24	0.98	2.33	
Türkiye	Overall	15.50	-	29.58	0.27	1.14	2.42	Soykan et al. <sup>39</sup>
Algeria	Female	21.75	-	34.13	0.26	1.50	2.48	Kherraz et al. <sup>41</sup>
	Male	17.50	-	26.78	0.38	0.75	2.43	
Egypt	Female	13.81	27.12	-	-	-	-	Azab et al. <sup>20</sup>
	Male	13.40	24.40	-	-	-	-	
	Overall	13.60	25.77	30.65	0.28	0.16	2.42	
Algeria	Overall	20.50	140.81	32.03	0.28	0.58	2.45	Dahel et al. <sup>21</sup>
	Female	16.77	48.84	37.14	0.15	1.33	2.32	
Türkiye	Male	16.72	48.25	29.95	0.22	0.95	2.29	This study
	Overall	16.74	48.50	37.50	0.15	1.32	2.33	

between 1.02 to 2.26  $y^{-1}$ ; 0.15 to 0.67  $y^{-1}$ ; 0.66 to 1.59  $y^{-1}$  and 0.46 to 0.87, respectively.<sup>4,7,20,21,38,39</sup> Our estimate  $E$  was 0.68 and higher than 0.50. Gulland<sup>30</sup> suggested that a fish stock is optimally exploited at a level of  $F$  which generates  $E = 0.50$ ,  $F_{opt} = M$ . Based on these suggestions, our results mean that the available stock is being heavily exploited.

Generally, many Sparidae species are long-lived fishes and have a low value of natural mortality; for instance, the natural mortality of *Pagrus pagrus*<sup>42</sup> and *Pagellus erythrinus*<sup>43</sup> were 0.22 and 0.43. In the bogue, the natural mortality was between 0.15  $y^{-1}$  and 0.67  $y^{-1}$ . While Pauly<sup>29</sup> estimated the natural mortality of fishes via von Bertalanffy growth parameters and assumed that there is a relation between fish size and natural mortality. The optimum exploitation rate ( $E$ ) is assumed to be close to 0.50.<sup>30</sup>  $E$  of bogue was 0.68 showing the stock at a higher optimum level of exploitation. Similar to these results, in Algarve, Algerian, Turkey, and Egypt waters.

High rates of fishing mortality (0.76) impact the population structure and growth rates of many species in addition to environmental considerations. Fishing generally alters the length structure of fishing stocks. In fact, *B. boops* catch increased in the waters of Türkiye in 2016, 2017, and 2018 (2795.1, 3175.0, and 3559.3 tones).<sup>2</sup> Other species, like bogue, have also shown this link between growth and exploitation. The relationship between fishing mortality and growth performance may ultimately be due to commercial fishermen's adaptation to changing fishing practices and fishing gear selectivity, which has the effect of continuously removing larger individuals from populations.

This study investigated the age, growth, and mortality of bogue, *B. boops* (Linnaeus, 1758) and provides basic biological data that are useful in sustainable fishery management of this species in Turkey waters (Antalya Bay, northeastern Mediterranean Sea).

#### ACKNOWLEDGMENTS

I am pleased to thank Prof. Dr. Seval BAHADIR KOCA (Is-parta) for improving the language and proofreading the manuscript.

#### DECLARATION OF COMPETING INTERESTS

The author declare that he has no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper.

#### DATA AVAILABILITY STATEMENT

The data that support the findings of this study are available from the corresponding author upon reasonable request.

#### ETHICS APPROVAL

No ethical approval was the need for this study. Fish specimens came from commercial fisheries.



AUTHORS' CONTRIBUTION - [CREDIT](#)

Submitted: June 01, 2023 CST. Accepted: July 02, 2023 CST.

Published: July 21, 2023 CST.

*Conceptualization, Investigation, Data curation, Methodology, Formal Analysis, Writing – original draft, and Writing – review & editing:* Habil Uğur Koca



This is an open-access article distributed under the terms of the Creative Commons Attribution 4.0 International License (CCBY-4.0). View this license's legal deed at <http://creativecommons.org/licenses/by/4.0> and legal code at <http://creativecommons.org/licenses/by/4.0/legalcode> for more information.

## REFERENCES

1. Bauchot ML, Hureau JC. Sparidae. In: Whitehead PJP, Bauchot ML, Hureau JC, Nielsen J, Tortonese E, eds. *Fishes of the Northeastern Atlantic and the Mediterranean*. Paris; 1986. doi:[10.2307/1444931](https://doi.org/10.2307/1444931)
2. TÜİK. *Turkish Fishery Statistics*. Turkish Statistical Institute; 2022.
3. Gordo LS. On the age and growth of bogue, *Boops boops* (L.), from the Portuguese coast. *Fish Man Eco*. 1996;3(2):157-164. doi:[10.1111/j.1365-2400.1996.tb00139.x](https://doi.org/10.1111/j.1365-2400.1996.tb00139.x)
4. Allam SM. Growth, mortality, and yield per recruit of bogue, *Boops boops* (L.), from the Egyptian Mediterranean waters off Alexandria. *Mediterr Mar Sci*. 2003;4(1):87-96. doi:[10.12681/mms.244](https://doi.org/10.12681/mms.244)
5. El-Haweet A, Hegazy M, Abu-Hatab H, Sabry E. Validation of length frequency analysis for *Boops boops* (Bogue) growth estimation. *Egypt J Aqua Res*. 2005;31(1):399-408.
6. Khemiri S, Gaamour A, Zylberberg L, Meunier F, Romidhane MS. Age and growth of bogue, *Boops boops*, in Tunisian waters. *Acta Adriatica*. 2005;46(2):159-175.
7. Monteiro P, Bentes L, Coelho R, et al. Age and growth, mortality, reproduction and relative yield per recruit of the bogue, *Boops boops* Linne, 1758 (Sparidae), from the Algarve (South of Portugal) logline fishery. *J Appl Ichth*. 2006;22:345-352. doi:[10.1111/j.1439-0426-2006.00756.x](https://doi.org/10.1111/j.1439-0426-2006.00756.x)
8. Karakulak FS, Erk H, Bilgin B. Length-weight relationships for 47 coastal fish species from the northern Aegean Sea, Turkey. *J Appl Ichth*. 2006;22:274-278. doi:[10.1111/j.1439-0426-2006.00736.x](https://doi.org/10.1111/j.1439-0426-2006.00736.x)
9. Özaydın O, Taşkavak E. Length-weight relationships for 47 fish species from İzmir Bay (eastern Aegean Sea, Turkey). *Acta Adria*. 2006;47:211-216.
10. Matasin Z, Vucnic S. *Ceratothoa oestroides* (Risso, 1826) in bogue (*Boops boops* L.) and picarel (*Spicara smaris* L.) from the Velebit channel in the Northern Adriatic. *Veterinarski Arhiv*. 2008;78(4):363-367.
11. El-Okda NL. Age and growth of *Boops boops* (L.) from Egyptian Mediterranean waters off Alexandria Egypt. *Egypt J Aqua Bio Fish*. 2008;12(1):13-23. doi:[10.21608/ejabf.2008.1968](https://doi.org/10.21608/ejabf.2008.1968)
12. Kara A, Bayhan B. Length-weight and length-length relationships of the bogue, *Boops boops* (Linnaeus, 1758) in İzmir Bay (Aegean Sea of Turkey). *Belg J Zoo*. 2008;138(2):154-157.
13. Kara A, Bayhan B. Age and growth of *Boops boops* (Linnaeus, 1758) in Izmir Bay, Aegean Sea, Turkey. *J Appl Ichthyol*. 2015;31(4):620-626. doi:[10.1111/jai.12680](https://doi.org/10.1111/jai.12680)
14. Kasalica O, Regner S, Durovic M. Some aspects of the biology of the bogue, *Boops boops* (Linnaeus, 1758) in Montenegrin waters (South Adriatic Sea). *Studia Marina*. 2011;25(1):59-72.
15. Bolognini L, Domenichetti F, Grati F, Polidori P, Scarcella G, Fabi G. Weight-length relationships for 20 fish species in the Adriatic Sea. *Turk J Fish Aqua Sci*. 2013;13(3):555-560. doi:[10.4194/1303-2712-v13\\_3\\_21](https://doi.org/10.4194/1303-2712-v13_3_21)
16. Özvarol Y. Length-weight relationships of 14 fish species from the Gulf of Antalya (northeastern Mediterranean Sea, Turkey). *Turk J Zoo*. 2014;38:342-346. doi:[10.3906/zoo-1308-44](https://doi.org/10.3906/zoo-1308-44)
17. Bottari T, Micale V, Liguori N, et al. The reproductive biology of *Boops boops* (Linnaeus, 1758) (Teleostei: Sparidae) in the southern Tyrrhenian Sea (central Mediterranean). *Cah Bio Mar*. 2014;55:281-292.
18. Layachi M, Idrissi MH, Ramdani M, Sahnouni S, Flower R. Growth and reproduction of the bogue *Boops boops* L., 1758 in the Mediterranean coastal area between Nador and Sidia (Morocco). *Bul L'Inst Sci, Rabat, Section Sciences de la Vie*. 2015;37:53-59.
19. Ceyhan T, Ertosluk O, Akyol O, Özgül A. The maximum size of Bogue, *Boops boops* (Perciformes: Sparidae) for the Mediterranean. *Acta Aqua Turcica*. 2018;14(4):399-403. doi:[10.22392/egirdir.463612](https://doi.org/10.22392/egirdir.463612)
20. M. Azab A, M. El-Far A, M. El-Sayed A. Age, growth and population structure of bogue, *Boops boops*, in the mediterranean waters front Alexandria, Egypt. *Egypt J Aqu Bio and Fish*. 2019;23(3):69-81. doi:[10.21608/ejabf.2019.35327](https://doi.org/10.21608/ejabf.2019.35327)
21. T. Dahel A, Rachedi M, Tahri M, Benchikh N, Diaf A, Djebar AB. Fisheries status of the bogue *Boops boops* (Linnaeus, 1758) in Algerian East Coast (Western Mediterranean Sea). *Egypt J Aqua Bio Fish*. 2019;23(4):577-589. doi:[10.21608/ejabf.2019.60554](https://doi.org/10.21608/ejabf.2019.60554)



22. Cengiz Ö, Paruğ ŞŞ, Kızılkaya B. Weight-length relationship and reproduction of Bogue (*Boops boops* Linnaeus, 1758) in Saros Bay (Northern Aegean Sea, Turkey). *KSU J Agri Natu*. 2019;22(4):577-582. doi:[10.18016/ksutarimdogavi.516700](https://doi.org/10.18016/ksutarimdogavi.516700)
23. Gayanilo FC, Sparre P, Pauly D. *F.A.O. ICLARM Stock Assessment Tools II (FiSAT II). User's Guide*. Revised version. F.A.O.; 2005.
24. Ricker WE. Computation and interpretation of biological statistics of fish populations. *Bul Fish Res Board Can*. 1975;191:1-382.
25. Le Cren ED. The length-weight relationship and seasonal cycle in gonad weight and condition in the Perch (*P. fluviatilis*). *J Ani Eco*. 1951;20(2):201-219. doi:[10.2307/1540](https://doi.org/10.2307/1540)
26. Sparre P, Venema SC. *Introduction to Tropical Fish Stock Assessment, Part 1*. F.A.O. Fisheries Technical Paper; 1992.
27. Pauly D, Munro JL. Once more on the comparison of growth in fish and invertebrates. *Fishbyte*. 1984;2(1):21.
28. Pauly D. *Fish Population Dynamics in Tropical Waters: A Manual for Use with Programmable Calculators*. ICLARM Studies and Reviews; 1984.
29. Pauly D. On the interrelationships between natural mortality, growth parameters, and mean environmental temperature in 175 fish stocks. *ICES J Mar Sci*. 1980;39(2):175-192. doi:[10.1093/icesims/39.2.175](https://doi.org/10.1093/icesims/39.2.175)
30. Gulland JA. *The Fish Resources of the Ocean*. Fishing News (Books) Ltd., West Byfleet; 1971.
31. Zar JH. *Biostatistical Analysis*. Third. Prentice-Hall; 1996.
32. Boudinar AS, Chaoui L, Kara MH. Age, growth and reproduction of the sand smelt *Atherina boyeri* Risso, 1810 in Mellah Lagoon (Eastern Algeria). *J Appl Ichth*. 2016;32(2):302-309. doi:[10.1111/jai.12992](https://doi.org/10.1111/jai.12992)
33. Froese R. Cube law, condition factor, and weight-length relationships: History, meta-analysis, and recommendations. *J Appl Ichth*. 2006;22(4):241-253. doi:[10.1111/j.1439-0426.2006.00805.x](https://doi.org/10.1111/j.1439-0426.2006.00805.x)
34. Allen KR. Application of the Bertalanffy growth equation to problems of fisheries management: A review. *J Fish Board Can*. 1969;26(9):2267-2281. doi:[10.1139/f69-222](https://doi.org/10.1139/f69-222)
35. Moutopoulos DK, Stergiou KI. Length-weight and length-length relationships of fish species from the Aegean Sea (Greece). *J Appl Ichth*. 2002;18:200-203. doi:[10.1046/j.1439-0426-2002-00281.x](https://doi.org/10.1046/j.1439-0426-2002-00281.x)
36. Frost WE. The age and growth of eels (*Anguilla anguilla*) from the Windemere catchment area, Part 2. *J Anim Ecol*. 1945;14(2):106-124. doi:[10.2307/1387](https://doi.org/10.2307/1387)
37. Bagenal TB, Tesch FW. Age and growth. In: Bagenal TB, ed. *Methods for Assessment of Fish Production in Fresh Waters*. IBP Handbook. Blackwell Scientific Publications; 1978:101-136.
38. Benia R, Mouffok S, Boutiba Z. Estimation of the exploitable biomass and the reference biological point, F0.1, of bogue *Boops boops* L., in the bay of Bou-Ismaïl, center Algerian. *J Biodiv and Environ Sci (JBES)*. 2014;5(2):420-427.
39. Soykan O, İlkyaz AT, Metin G, Kınacıgil HT. rowth and reproduction of *Boops boops*, *Dentex macrophthalmus*, *Diplodus vulgaris*, and *Pagellus acarne* (Actinopterygii: Perciformes: Sparidae) from East-Central Aegean Sea, Turkey. *Acta Ichth Pisc*. 2015;45(1):39-55. doi:[10.3750/aip2015.45.1.05](https://doi.org/10.3750/aip2015.45.1.05)
40. Bouazizi A, Djebbour O, Benia R. Estimation de la biomasse exploitable et du point de reference biologique, F0.1, de la bogue de la region d'Alger, Congres International Gestion systemique des ressources halieutiques. *GSRH*. Published online 2010:29-30.
41. Kherraz A, Kherraz A, Boutiba Z. Interrelationship age and growth of *Boops boops* (Linnaeus, 1758) in Western Mediterranean coasts of Algeria. *Adv Environ Bio*. 2016;10(4):140-145.
42. Vassilopoulou V, Papaconstantinou C. Age, growth and mortality of the Red Porgy, *Pagrus pagrus*, in the Eastern Mediterranean Sea (Dodecanese, Grece). *Vie Milieu*. 1992;42(1):51-53.
43. Gurbet R, Akyol O, Yalçın E. Catch per unit effort and mortality rates of two spraid species, *Pagellus acarne* and *Pagellus erythrinus* from bottom trawl fishery in İzmir Bay, Aegean Sea. *J Ani Vet Adv*. 2012;11(5):681-686. doi:[10.3923/javaa.2012.681.686](https://doi.org/10.3923/javaa.2012.681.686)