

## Original Research Articles

# Transcriptome analysis reveals the important roles of a two-component system, flagellar assembly, active efflux system and outer membrane proteins in the anti-quinolone ability of *Vibrio harveyi* from orange-spotted grouper (*Epinephelus coioides*)

Xiaochen Tang<sup>1</sup>, Yu Ding<sup>1a</sup>

<sup>1</sup> Guangdong Provincial Key Laboratory of Aquatic Animal Disease Control and Healthy Culture, Southern Marine Science and Engineering Guangdong Laboratory (Zhanjiang), Guangdong Key Laboratory of Control for Disease of Aquatic Economic Animals, Fisheries College of Guangdong Ocean University, Zhanjiang, China

Keywords: flagellar assembly, quinolone-resistance, transcriptome, two-component system, *Vibrio harveyi*

<https://doi.org/10.46989/001c.84647>

## Israeli Journal of Aquaculture - Bamidgheh

Vol. 75, Issue 2, 2023

The drug resistance of *Vibrio harveyi* in aquaculture became more severe because the control of Vibriosis depends majorly on the current antibiotics. Transcriptomes of a wild-type strain (VS) and its quinolone-resistant mutants (VR) of *V. harveyi* were respectively sequenced by RNA-seq technology. A total of 2,082 unigenes were obtained after *de novo* splicing and assembly. 129 genes were identified with significant differential expression in strain VR compared to strain VS, among which 65 were up-regulated and 64 down-regulated. Then, functional annotation and enrichment analysis of these differentially expressed genes (DEGs) were performed. GO enrichment results showed that DEGs focused mainly on cell structure, substance metabolism, and transporter. COG classification of the DEGs mainly focused on amino acid transport and metabolism, cell wall/membrane biosynthesis, carbohydrate transport and metabolism, ribosomal structure, and biosynthesis. KEGG pathways related to a two-component system, ABC transport system and flagellar assembly (ko02040) were enriched significantly, and 9 genes associated with quinolone-resistance ability, including genes for resistance-related transport proteins, outer membrane proteins, and DNA repair-related proteins were discovered through analysis of the drug-resistance related genes. Ten DEGs (including the above part genes of 9 drug resistance-related genes) in the transcriptome data were taken to analyze their expression with real-time qPCR. The results were the same as the changes of the above transcriptome analysis, further confirming the reliability of the transcriptome sequencing and data analysis. In a word, genes from a two-component system, flagellar assembly, active efflux system and outer membrane proteins take great roles in the quinolone-resistance of *V. harveyi*. These results provide enough information for further study on the molecular mechanism of quinolone-resistance and give a helpful transcriptomic resource to unravel the contact between quinolone-resistance and metabolic pathways in *Vibrios*.

## INTRODUCTION

*Vibrio harveyi* is widely distributed in the marine environment and many marine organisms, and it is an opportunistic pathogen of a variety of marine fish and invertebrates such as the pathogen of luminous vibriosis in the farmed penaeid prawns.<sup>1-3</sup> Though the established methods of infectious disease prevention, such as immunization with vaccines, have been investigated and developed in the past ten years, control of *V. harveyi* in aquaculture depends ma-

jorly on antibiotics at present.<sup>4-8</sup> But it is well known that antibiotic compounds are associated with various problems and side effects on humans and the environment.<sup>9</sup> Drug resistance of bacteria in aquaculture became more serious.<sup>9-11</sup>

With the development of high-throughput sequencing technology, next-generation sequencing (NGS) technologies have improved the efficiency and speed of gene discovery and its function study with high throughput and low expenditure.<sup>12</sup> RNA sequencing (RNA-Seq) can detect

a Corresponding author: Yu Ding. Tel.: +86-759-2383509, fax: +86-759-2383507, e-mail: dingyuddd@163.com

the expression level of thousands of genes simultaneously and deeply analyze the functional pathways and regulatory methods in biological processes.<sup>13</sup> It can establish expression profiling and compare expression patterns between different strains or stages from the same organisms. So the technology has been widely used to discover potential drug-related genes, such as drug-resistant genes.<sup>13</sup>

Quinolone is a synthetic broad-spectrum antibacterial drug with a basic structure of the 4-quinolone ring. For example, because of their high antibacterial activity, norfloxacin and enrofloxacin are widely used to prevent and treat bacterial diseases in various aquatic animals. The resistance of bacteria to quinolone drugs in the aquaculture environment is becoming increasingly apparent.<sup>11,14</sup> The drug-resistant mechanism of bacteria to quinolones is mainly relative to the mutation-mediated changes of the topoisomerase target site (QRDRs, Quinolone Resistance Determining Regions, specific site), plasmid (coding drug-resistance protein and efflux pump), chromosome (coding porin and efflux pump) and so on.<sup>15</sup>

At present, the resistance mechanism of quinolone has been studied on aquatic pathogens such as *V. parahaemolyticus*,<sup>16</sup> *Aeromonas hydrophila*,<sup>17</sup> and *V. cholerae*.<sup>14</sup> QRDRs-specific site mutation and plasmid-mediated quinolone resistance genes (PMQR) *qnrVC5* have been found and researched in *V. parahaemolyticus*.<sup>16</sup> There were single-point mutations on GyrA and ParC in *V. parahaemolyticus*, which was highly resistant to ciprofloxacin. It is suggested that the mutation of the target site is an essential mechanism of drug resistance in *V. parahaemolyticus*.<sup>16</sup> *qnrVC5* have been found on the plasmids of four *V. parahaemolyticus* strains.<sup>16</sup> In *A. hydrophila*, double mutations of QRDRs on *gyrA* and *parC* have also been reported, and the strains with double mutations show a high drug resistance.<sup>17</sup> But there is a lack of research on the resistance mechanism to quinolone in *V. harveyi*, an important marine aquatic pathogen in China.<sup>3</sup> In this study, transcriptome sequencing was performed on the resistant and sensitive strains of *V. harveyi* to obtain the changes in the transcriptome level and to disclose the relation between its drug resistance and metabolic pathway. The results can provide important data for further exploration of the molecular mechanism of quinolone resistance in *V. harveyi*.

## MATERIALS AND METHODS

### BACTERIA

*V. harveyi* wild-type strain (VS), which is sensitive to quinolone (MIC of 0.5 µg/ml), was isolated from diseased orange-spotted grouper clinically in Haikou (Hainan Province, China) and deposited at -80 °C in our laboratory, Guangdong Provincial Key Laboratory of Aquatic Animal Disease Control and Healthy Culture in China. The VS strain of *V. harveyi* was used to induce the drug-resistant mutant in stepwise exposure to sub-inhibitory with gradually increasing concentrations of norfloxacin (0.125, 0.25, 0.5 µg/ml, respectively) *in vitro*. The drug resistance and biological characteristics of strains before and after induc-

tion were detected to screen the drug-resistant mutants. Three biological replicates were performed. A high-level quinolone-resistant mutant (VR strain, MIC of 512 µg/ml) with stable resistance and cross-resistance was successfully constructed and deposited at -80 °C. Before using, bacteria strains VS and VR were cultured and activated in tryptic soy agar (TSA, pH7.5) containing 15 g/l tryptone, 5 g/l soy peptone, and 5 g/l NaCl at 28 °C for 18 h. Then, they were cultured in a TSB medium. Cells were collected for the following RNA extraction by centrifugation when OD<sub>600nm</sub> reached 0.5 (28 °C for 12 h).

### RNA LIBRARY CONSTRUCTION

Total RNA was extracted following the manufacturer's instructions using TransZol Up Plus RNA Kit (Transgen Company, Beijing) and DNA digestion with DNase I (TaKaRa, Japan). The quantity and quality of the RNA samples were measured by gel electrophoresis, Qubit 2.0, Agilent 2100, and NanoDrop 2000 (Thermo, USA). High-quality RNA of the exact quantities was used in the subsequent experiments. Three biological replicates of each group (VR and VS) were used to construct cDNA libraries.

The cDNA libraries were constructed following the instructions for the Illumina Gene Expression Sample Prep Kit (Illumina Inc., San Diego, CA, USA). In detail, the total mRNA was mixed with the fragmentation buffer and fragmented into short fragments, which were used as templates for cDNA synthesis. First-strand cDNA synthesis was carried out by random hexamer primer, and M-MuLV reverse transcriptase, and the second-strand cDNA was synthesized by DNA polymerase I and RNase H. The cDNA fragments were purified and resolved with EB buffer for end reparation and single nucleotide A (adenine) addition, and then, they were connected with adapters. The adaptor-modified fragments were selected by gel purification and amplified through PCR to create the final cDNA library.

Use Qubit 2.0 to detect the concentration of the library, and use qPCR to determine the effective concentration of the library for the library quality control. Then, high-throughput sequencing was performed on the HiSeq2500 platform (Illumina Inc., San Diego, CA, USA). Three sequencing runs used different kits, and all samples were run the same.

### BIOINFORMATICS ANALYSIS

Clean Data, which came from raw data after removing the connector sequence and unknown and low-quality reads to avoid negatively affecting the bioinformatics analysis, were assembled to obtain the unigenes library of the species. The transcripts' structure was analyzed, and functional annotation and enrichment analysis of the differentially expressed genes (DEGs) were performed. Sequence assembly and annotation were listed in detail as follows.

*De novo* transcriptome assembly was performed with Trinity software.<sup>18</sup> The resulting sequences obtained from Trinity were called unigenes.

The unigenes were compared against diverse protein databases, including the nonredundant protein (Nr) data-

base (NCBI) and Swissprot databases (<http://www.uniprot.org/blast/>). The results of annotation from each protein database were presented independently. Blast2GO (<http://www.blast2go.com/b2ghome>) was used to obtain GO annotation of the unigenes.<sup>19</sup> The COG database (<ftp://ftp.ncbi.nih.gov/pub/COG/COG/>) was also used to predict and classify functions of the unigenes.

#### IDENTIFICATION OF DGES

Gene expression levels were calculated as unit fragments per kilobase per million reads (FPKM), and the correlation between the two groups of samples was calculated. EBSeq software was used for differential expression analysis, a stringent algorithm p-value was used to identify DEGs between VR and VS strains according to the study,<sup>20</sup> and Benjamini-Hochberg method was used to correct the significant p-value (p-value).<sup>21</sup> The false discovery rate (FDR) was used to determine the p-value threshold in multiple tests to judge the significance of the gene expression differences.  $FDR \leq 0.05$  and the absolute value of fold-change ( $\geq 2$ ) were used as the default screening conditions to assess DEGs in this study. For GO and KEGG enrichment analyses, all differentially expressed genes were mapped to terms in GO and the KEGG database by a hypergeometric test using the whole transcriptome as the background as above.

#### QRT-PCR

In this study, we randomly selected 10 unigenes from the sequencing results (including resistance-related genes, *recA*, *recN*, *ompU*, *acrB* and *ompA*, and others, *mfp*, *V0780*, *V1668*, *mepA* and *ompV*) for the real-time fluorescence quantification PCR analysis (qRT-PCR) to validate the results of differential expression analysis. The sequencing results showed that fluorescent quantitative primers were designed using Primer5.0 ([Table 1](#)). cDNA was synthesized with Easy Script One-step gDNA Removal and cDNA Synthesis SuperMix in a total volume 20  $\mu$ L (Total RNA (400ng/ $\mu$ L) 5 $\mu$ L, gDNA Remover 1 $\mu$ L, Random primer 1 $\mu$ L, 2 $\times$ ES Reaction Mix 10 $\mu$ L, Easy Script RT Enzyme Mix 1 $\mu$ L, and RNase-free water 2 $\mu$ L). Firstly, the samples were incubated at 25 °C for 10 minutes, followed by a 30-minute incubation at 42°C. Finally, the samples were incubated at 85 °C for 1 minute to inactivate the enzyme. Real-time qPCR was performed with SYBR® Premix Ex Taq™ (TaKaRa, Japan) on an iQ5 Real-Time PCR system (Bio-Rad, USA) in 20  $\mu$ L system (Forward Primer 0.5  $\mu$ L, Reverse Primer 0.5  $\mu$ L, 2 $\times$ TransStart Tip Green qPCR SuperMix 10  $\mu$ L, cDNA 2  $\mu$ L and dd H<sub>2</sub>O 7  $\mu$ L) under the following conditions: denaturation at 50 °C for 120 s and 95 °C for 120 s, then 40 cycles of 94 °C for 15 s, 60 °C for 15 s, and 72 °C for 60 s. Relative gene expression was analyzed by the  $2^{-\Delta\Delta C_t}$  method.<sup>22</sup> *gapA* was chosen as the internal control for normalization. Samples were assessed in triplicate. SPSS 17.0 software was used to determine statistical significance and draw plots with Excel 2010.

## RESULTS

### SEQUENCE ANALYSIS AND DE NOVO ASSEMBLY

Compared with VS, VR's swimming ability and biofilm formation with a high-level quinolone resistance significantly increased, and the growth curve had no noticeable changes (Data not shown). The quality (concentration and RNA Integrity Number (RIN)) of RNA extracted from the above two strains is high enough to meet the requirements for the next library construction. The sequencing and assembly results are shown in **Table S1** in the supplementary material, which contains 56,440,529 raw reads. Strain VS obtained 29,171,777 clean reads (91.0 % of the original data), and strain VR obtained 23,371,075 (95.9 % of the original data). Q30 base percentages are at least 88.41 %. These showed that high-quality clean data was obtained, ensuring the accuracy of sequence assembly and subsequent analysis. FPKM value (closer to 1) also indicated a better reproducibility of the two groups.

The total transcriptome data (NCBI accession number PRJNA791968) was spliced *de novo*, and then assembled and get a total of 2,082 unigenes with a total length of 4,687,723 bp and an average length of 2,251 bp. The number with a length between 200 and 300 bp is the largest, in total 745 and accounting for 35.78 %. And the unigenes with more than 3,000 bp is 379, accounting for 18.2 % (**Figure S1** in the supplementary material).

### SEQUENCE ANNOTATION

Using Glimmer soft to perform gene prediction on the uni-gene library, a total of 2,519 genes and their CDS sequences were obtained, and the average length of the predicted genes was 1,190 bp. The predicted genes were annotated functionally, and finally obtain 2,509 genes with the annotated information (**Table S2** in the supplementary material).

### DIFFERENTIALLY EXPRESSED GENES

129 DEGs were found finally. Compared with strain VS, drug-resistant mutant VR had 65 genes up-regulated and 64 genes down-regulated, of which 5 genes were only expressed in drug-resistant bacteria. The volcano diagram of DEGs was shown in **Figure 1A**. DEG correlation scatter plot was shown in **Figure 1B**.

### ENRICHMENT ANALYSIS OF DEGS

DEGs between strain VS and VR and the annotated results of all genes at the GO secondary node are shown in **Figure 2**. The results showed that all unigenes are annotated into 38 second-level entries, of which DEGs are annotated into 26. Regarding molecular functions, they mainly focus on transporter activity and binding. For cell composition, major categories with differences were cell (includes the plasma membrane and any external encapsulating structures such as the cell wall and cell envelope), membrane, and cell part (any constituent part of a cell). Among biolog-

**Table 1. Primers for RT-PCR.**

Gene	Strand	Primers (5'-3')	Product (bp)
<i>acrB</i>	F	GACGGGTGAGTTTATGGGC	222
	R	AACCAACGGAAACGAAGTG	
<i>mfp</i>	F	CACCAGTGAGCCAAATGCG	125
	R	TAATGCCTGCCGCTACCA	
V0780	F	AAGTGAAGCAGGGCGATGT	200
	R	TTGCTCAAGGTTTGCCAGT	
V1668	F	GAGATGGTGGTTCAGTCGC	200
	R	GTTACCGCTTTCGTTGTGAG	
<i>mepA</i>	F	CCAGCCAGATGTGGTTTAGC	190
	R	ACCCAAGCGAGGGAATAGA	
<i>recA</i>	F	AGTGCCTCAAGAGCGGTGTTT	143
	R	CAAGCGGTTTGTATATTAGGTT	
<i>recN</i>	F	AACAGCCAGCAAAATCAGG	145
	R	AGCTAGCGGCAAGTTCACC	
<i>ompU</i>	F	AGGATAGCCAAGATAAATACGG	135
	R	AGTAAGCGATACCACCAAGGA	
<i>ompA</i>	F	GGTGATGGCTTGTTGCG	147
	R	GCCCTTGCTGTCGGTATGA	
<i>ompV</i>	F	GGCTTCCAAGGTGGTGCTA	156
	R	CTTTGTATGCTTTGCGGTTT	
<i>gapA</i>	F	ATTAGATCCTCCACCTTGCTG	149
	R	TGTGAGAACAGAACTTTATGAGCC	

ical processes, metabolic processes and cellular processes were the most represented. The results show that drug-resistant bacteria differ from sensitive bacteria in cell structure, substance metabolism, and transport.

The function of these DEGs was predicted by comparing them with COG databases. The statistical results of COG classification of the DEGs are shown in [Figure 3](#). 90 DEGs were mapped to 19 different functional categories, which mainly focused on amino acid transport and metabolism, cell wall/membrane biosynthesis, carbohydrate transport and metabolism, ribosomal structure, and biosynthesis.

The enrichment degree of KEGG pathway was analyzed using the enrichment factor, and Fisher's test was used to calculate the significance of enrichment. 78 DEGs were annotated to 34 metabolic pathways. KEGG classification map is shown in [Figure 4](#), and [Table 2](#) shows the enriched top 20 channels. The results showed that the pathway two-component system (ko02020), ABC transport system (ko02010), and flagella assembly (ko02040) have the largest number of the annotated differential genes, and they are enriched significantly ([Figure S2-S4](#) in the supplementary material).

#### ANALYSIS OF THE RESISTANCE-RELATED GENES

Some genes related to quinolone resistance were identified from the DEGs, including resistance-related transport proteins, outer membrane proteins, and DNA repair-related


proteins ([Table 3](#)). The results showed that 5 drug resistance transporter genes were up-regulated, among which resistance protein B (BSP001668) was up-regulated most. Genes of two outer membrane proteins (OmpA and OmpU) are down-regulated. The expression levels of DNA repair protein RecN and DNA recombinant protein RecA are both up-regulated. Since the target of quinolones is topoisomerase in cells, proteins related to DNA repair may play an indirect role in its drug resistance.

#### RESULTS WITH FLUORESCENCE QUANTIFICATION PCR

Ten genes were selected from the transcriptome data, and their fluorescence quantitative analysis was performed ([Figure 5](#)). The expression level of 8 genes, except genes *ompA* and *ompU* increased differently. Of which, gene *V1668* and *recN* increased with the most degree. The fold changes of *fmp*, *V0780*, and *recA* are also more than 1. The results showed that the expression changes of these genes are consistent with the transcriptome analysis results. Overall, the transcriptome sequencing data in this study is reliable.

#### DISCUSSION

The resistance mechanism of *Vibrio* to antibiotics is very complex.<sup>15</sup> The single-drug induction method adopted in this study can exclude the interference of the other en-



**Figure 1. Volcano plot and correlation scatter plot of the differentially expressed genes of the wild-type strain (VS) and its quinolone-resistant mutants (VR)**


Note: Volcano plot of the differentially expressed genes of VR and VS (A), 65 genes up-regulated and 64 genes down-regulated in drug-resistant strain VR; Differentially expressed gene correlation scatter plot (B).

vironmental factors in the bacteria resistance. So the obtained drug-resistant strain VR can be used as a single sample induced by quinolone.

The next-generation high-throughput sequencing technology RNA-Seq was used for transcriptome sequencing analysis of quinolone-sensitive and induced drug-resistant strains of *V. harveyi* in the study. A total of 13.23 Gb Clean Data was obtained, providing sufficient and adequate data and information for the following research (Table S1 in the supplementary material). The Assembly *de novo* technique was used to obtain the transcripts of the species, and a uni-


gene library of the species was constructed, which obtained a total of 2,082 unigenes (Figure S1 in the supplementary material). In this study, there were 129 differentially expressed genes, 65 of which were up-regulated and 64 were down-regulated in mutant strain VR (Figure 1). But Li et al.<sup>24</sup> analyzed the transcriptome of ciprofloxacin-induced *Salmonella typhimurium* ATCC13311-resistant strains. They obtained 3,466 differentially expressed genes compared with the sensitive bacteria, of which 3,434 were up-regulated. The down-regulated genes had only 32 genes. This result may be due to the differences in species and to the






**Figure 2. GO categories of unigenes and the differentially expressed genes (DEGs)**

Note: Blast2GO (<http://www.blast2go.com/b2gohome>) was used to obtain GO annotation of the unigenes. The unigenes were functionally classified under three main functional categories: cellular component, molecular function and biological process. The right y-axis indicates the number of unigenes in each subcategory. The left y-axis indicates the percentage of a specific category of unigenes in that main category.



**Figure 3. COG function classification of the differentially expressed genes (DEGs)**

Note: The function of the differentially expressed genes was predicted by comparing DEGs of the study with COG databases. 90 differentially expressed genes were mapped to 19 different functional categories.



**Figure 4. KEGG classification map.**

different drugs. It also showed that it is necessary to study further the action mechanism of drugs and the drug resistance mechanism of the different bacteria.

The annotations of unigenes can provide gene functions, physiological processes, and pathways involved in *Vibrio*.<sup>13</sup> Combining the functional annotation and enrichment analysis results of differentially expressed genes and the resistance-related genes analysis result, DEGs of the two-component system (TCS), flagella assembly, active efflux system, outer membrane proteins, DNA repair system, and cell membrane structure were discovered in quinolone-resistant *V. harveyi* (Figure 2-5). But QRDRs and *qnrVC5* have not been found, like in *V. Parahaemolyticus*.<sup>16</sup>

There are 12 differentially expressed genes annotated to the two-component system pathway (ko02020) (Figure 4 and Table 2). TCS is a signal transduction system for bacteria responding to environmental changes.<sup>25</sup> It regulated multiple functions of bacteria, including bacterial growth, biofilm-forming, drug resistance, virulence, quorum sensing, etc.<sup>26</sup> Among the up-regulated genes, it was found that the *flhC* (BSP000991) gene, relating to the assembly of flagellar (Table 2), was regulated by a two-component system. Flagella assembly plays a significant role in bacterial drug

resistance.<sup>25</sup> Transcriptome results showed that the expression of flagella-related proteins increased in the drug-resistant *Vibrio*. This is consistent with our previous research results that the phenotypic characteristics of drug-resistant bacteria, such as their mobility and biofilm formation ability, were increased (data not shown). QseBC, some TCS of *E. coli*, also participated in the transcriptional regulation of flagellar regulation-related genes (*flhD*, *flhC*, *flhA*, *motA*, *fliC*).<sup>26</sup> Flagellar expression increased in the present study, and as a result, the drug-resistant strain exhibited increased mobility and biofilm formation. So the regulation function of TCS on the flagella-related proteins is very important to drug-resistance of bacterium. This may be an indirect and non-specific drug-resistance mechanism for *V. harveyi* like that in *E. coli*.

Active efflux system is important in bacterial multi-drug resistance.<sup>27</sup> In our results, four drug resistance-related differentially expressed transporters, AcrB (BSP000778), RND family efflux transporter (BSP000779), resistance protein (BSP000780), and resistance protein B (BSP001668) belongs to the RND efflux pump family of active efflux system. Multidrug transporter MatE (BSP002444) belongs to the MATE family (Table 2 and Table 3). The RND-type ef-

**Table 2. Top 20 pathways with the greatest number of annotated sequences.<sup>1</sup>**

Pathway	Pathway ID	DGEs with pathway annotation	All genes with Pathway annotation
Ribosome	ko03010	4	7
Two-component system	ko02020	12	78
Phosphonate and phosphinate metabolism	ko00440	2	2
ABC transporters	ko02010	14	119
Butanoate metabolism	ko00650	5	30
Synthesis and degradation of ketone bodies	ko00072	2	5
Flagellar assembly	ko02040	4	33
Riboflavin metabolism	ko00740	1	3
Glyoxylate and dicarboxylate metabolism	ko00630	2	16
Folate biosynthesis	ko00790	1	6
Protein export	ko03060	1	6
Valine, leucine and isoleucine degradation	ko00280	2	20
Aminobenzoate degradation	ko00627	1	7
Alanine, aspartate and glutamate metabolism	ko00250	2	21
Glycine, serine and threonine metabolism	ko00260	2	22
One carbon pool by folate	ko00670	1	8
Benzoate degradation	ko00362	1	9
beta-Alanine metabolism	ko00410	1	9
Arginine and proline metabolism	ko00330	2	25

<sup>1</sup>Pathway assignments conducted based on the KEGG database (<http://www.genome.jp/tools/blast/>), which contains graphical representation of biological processes.<sup>23</sup>

**Table 3. Differential expression of the candidate resistance-related gene.<sup>1</sup>**

ID	Gene length	log2FC	regulated	Swissprot annotation
BSP000778	3060	2.426	up	multidrug transporter AcrB
BSP000779	1062	2.674	up	RND family efflux transporter
BSP000780	1074	2.8512	up	resistance protein
BSP001668	1329	5.1206	up	resistance protein B
BSP002444	1353	1.8916	up	multidrug transporter MatE
BSP001484	681	-1.0375	down	Outer membrane protein A
BSP000701	969	-2.8633	down	Outer membrane protein U
BSP002256	1665	3.4331	up	DNA repair protein RecN
BSP001507	693	2.4831	up	DNA recombination protein RecA

<sup>1</sup> log2FC : log2 value of the times of the expression differences.


flux pump AcrAB-TolC system is an active efflux system in *E. coli*. Its related substrates include quinolones,  $\beta$ -lactams, macrolides, chloramphenicol, tetracycline and other antibiotics.<sup>28</sup> MATE transporter YdhE of *E. coli* can also transport quinolones and macrolide drugs.<sup>29</sup> *V. parahaemolyticus* NorM, mediating resistance to quinolone drugs, has 57 % identity and high similarity (88 %) with *E. coli* YdhE and belongs to the MATE family of transporters.<sup>27</sup> Our results showed that the resistance of *V. harveyi* to quinolone drugs is likely related to the efflux pump system.

The outer membrane's permeability also affects bacteria's sensitivity to quinolones.<sup>30</sup> It has been found that the outer membrane proteins of *E. coli* OmpF and OmpC are related to the diffusion of quinolone drugs. Hydrophilic

drugs (ciprofloxacin, norfloxacin) diffuse mainly into bacteria through OmpF.<sup>30</sup> The expression of two outer membrane proteins (OmpA and OmpU) decreased in our study (the relative expression level < 0) (Table 3), of which OmpA and *E. coli* OmpF have a high homology. Decreased expression of outer membrane proteins can reduce the diffusion of quinolone drugs and enable bacteria to obtain drug resistance. So outer membrane proteins (OmpA and OmpU) can be related to the quinolone resistance of *V. harveyi*.

Because real-time fluorescent quantitative PCR has high sensitivity and reproducibility, it can detect small differences in gene expression, and it can be used to measure the expression level of genes and verify the results of transcriptome analysis and their functions in drug resistance of





**Figure 5. The result of quantitative Real-time PCR.**

Note: Ten unigenes from the sequencing results (including 9 drug resistance-related genes) were selected for quantitative RT-qPCR to validate the results of differential expression analysis in transcriptome data.

*V. harveyi*.<sup>22</sup> In this study, the specific primers and the reaction system with good melting curve were selected. The housekeeping gene *gapA* in bacteria was selected as the internal reference gene for qPCR after the reaction conditions and system, and the primers were optimized. qPCR results are consistent with the transcriptome analysis such that the expression of *ompA* and *ompU* also decreased (the relative expression level < 1), and the expression of the other genes also increased (Figure 5), confirming the reliability of transcriptome sequencing and data analysis, and confirming that the above-related genes, *acrB*, *ompA*, *ompU*, *recA* and *recN*, may play key roles in the drug resistance of *V. harveyi*.

The metabolic pathways of two-component systems, flagella assembly, active efflux system, and outer membrane proteins, are significantly different, suggesting that changes in various bacterial metabolism pathways can lead to drug resistance. This is an important supplement to the study on drug resistance mechanisms though QRDRs and differentially expressed *qnrVC5* have not been found in *V. harveyi*. Results provide essential data for exploring the molecular mechanism of quinolone resistance in *V. harveyi* and present a great reference to unravel the mechanism of quinolone resistance in the other different *Vibrios*. The re-

lationship between the drug resistance and the metabolic pathway deserves further exploration.

## CONCLUSION

This study compared *V. harveyi* transcriptome profiles between the wild-type strain and its quinolone-resistant mutants. Comparative transcriptome analyses identified 9 genes associated with quinolone-resistance ability, some further verified by qPCR. The resistance mechanism of *V. harveyi* to quinolones is related to multiple pathways. The two-component system, flagella assembly, active efflux system, outer membrane protein, and DNA repair system constitute a complex network in developing drug resistance of *V. harveyi*, and their relation or co-operation between any two need to be studied further. Taken together, our results provide enough information for quinolone-resistance study of *V. harveyi*. And also provides a reference for the further research on the physiological changes and metabolic processes in the resistance formation of *Vibrios*.

## ACKNOWLEDGMENTS

Thank all the people who have dedicated time to the experiments. This work was supported by grants from the Natural Science Foundation of Guangdong Province (2022A1515012203) and the National Key Technology Support Program of China (No. 2012BAD17B03).

draft: Xiaochen Tang (Lead). Writing – review & editing: Xiaochen Tang (Equal), Yu Ding (Equal). Funding acquisition: Yu Ding (Lead). Resources: Yu Ding (Lead). Supervision: Yu Ding (Lead).

Submitted: April 27, 2023 CST. Accepted: July 15, 2023 CST.  
Published: August 07, 2023 CST.

## AUTHORS' CONTRIBUTIONS PER CREDIT

Conceptualization: Xiaochen Tang (Equal), Yu Ding (Equal).  
Methodology: Xiaochen Tang (Equal), Yu Ding (Equal). Formal Analysis: Xiaochen Tang (Lead). Writing – original



This is an open-access article distributed under the terms of the Creative Commons Attribution 4.0 International License (CCBY-4.0). View this license's legal deed at <http://creativecommons.org/licenses/by/4.0> and legal code at <http://creativecommons.org/licenses/by/4.0/legalcode> for more information.

## REFERENCES

1. Wang J, Ding Q, Yang Q, Fan H, Zhao P. *Vibrio alginolyticus* triggers inflammatory response in mouse peritoneal macrophages via activation of NLRP3 inflammasome. *Frontiers in Cellular and Infection Microbiology*. 2021;11:769777-769777. doi:[10.3389/fcimb.2021](https://doi.org/10.3389/fcimb.2021)
2. Zhou K, Tian KY, Liu XQ, et al. Characteristic and otopathogenic analysis of a *Vibrio alginolyticus* strain responsible for chronic Otitis Externa in China. *Front Microbiol*. 2021;12:750642-750642. doi:[10.3389/fmicb.2021.750642](https://doi.org/10.3389/fmicb.2021.750642). PMID:34975783
3. Mohi MM, Kuratani M, Miyazaki T, Yoshida T. Histopathological studies on *Vibrio harveyi*- infected tiger puffer, *Takifugu rubripes* (Temminck et Schlegel), cultured in Japan. *Journal of Fish Diseases*. 2010;33(10):833-840. doi:[10.1111/j.1365-2761.2010.01184.x](https://doi.org/10.1111/j.1365-2761.2010.01184.x)
4. Ragab RH, Elgendy MY, Sabry NM, et al. Mass kills in hatchery-reared European seabass (*Dicentrarchus labrax*) triggered by concomitant infections of *Amyloodinium ocellatum* and *Vibrio alginolyticus*. *International Journal of Veterinary Science and Medicine*. 2022;10(1):33-45. doi:[10.1080/23144599.2022.2070346](https://doi.org/10.1080/23144599.2022.2070346). PMID:35573137
5. Bedekar MK, Kole S. Fundamentals of fish vaccination. *Methods in Molecular Biology*. 2022;2411:147-173. doi:[10.1007/978-1-0716-1888-2\\_9](https://doi.org/10.1007/978-1-0716-1888-2_9)
6. Mohamad N, Mohd Roseli FA, Azmai MNA, et al. Natural concurrent infection of *Vibrio harveyi* and *V. alginolyticus* in cultured hybrid groupers in Malaysia. *J Aquat Anim Health*. 2019;31(1):88-96. doi:[10.1002/aah.10055](https://doi.org/10.1002/aah.10055)
7. Wang H, Zhu F, Huang Y, Ding Y, Jian J, Wu Z. Construction of glutathione peroxidase (GPx) DNA vaccine and its protective efficiency on the orange-spotted grouper (*Epinephelus coioides*) challenged with *Vibrio harveyi*. *Fish & Shellfish Immunology*. 2017;60:529-536. doi:[10.1016/j.fsi.2016.11.018](https://doi.org/10.1016/j.fsi.2016.11.018)
8. Sun BG, Dang W, Sun L, Hu YH. *Vibrio harveyi* Hsp70: Immunogenicity and application in the development of an experimental vaccine against *V. harveyi* and *Streptococcus iniae*. *Aquaculture*. 2014;418:144-147. doi:[10.1016/j.aquaculture.2013.10.018](https://doi.org/10.1016/j.aquaculture.2013.10.018)
9. Okuda J, Hayakawa E, Nishibuchi M, Nishino T. Sequence analysis of the *gyrA* and *parC* homologues of a wild-type strain of *Vibrio parahaemolyticus* and its fluoroquinolone-resistant mutants. *Antimicrob Agents Chemother*. 1999;43(5):1156-1162. doi:[10.1128/aac.43.5.1156](https://doi.org/10.1128/aac.43.5.1156). PMID:10223929
10. Zeng D, Feng J, Xu L, Su Y, Guo Z. Antibioigram types of *Vibrio harveyi* strains isolated from maricultured fish. *Journal of Fishery Sciences of China*. 2015;12:129-138.
11. Qiao Y, Wan XH, Shen SH. Research progress in antibacterial drugs resistance of Chinese aquaculture industry. *Chinese Journal of Antibiotics*. 2015;40:389-395.
12. Metzker ML. Applications of next-generation sequencing technologies—the next generation. *Nat Rev Genet*. 2009;11(1):31-46. doi:[10.1038/nrg2626](https://doi.org/10.1038/nrg2626)
13. Liu T, Zhu S, Tang Q, Chen P, Yu Y, Tang S. De novo assembly and characterization of transcriptome using Illumina paired-end sequencing and identification of Csa gene in ramie (*Boehmeria nivea* L. Gaud). *BMC Genomics*. 2013;14(1):125. doi:[10.1186/1471-2164-14-125](https://doi.org/10.1186/1471-2164-14-125). PMID:23442184
14. Po KHL, Wong MHY, Chen S. Identification and characterisation of a novel plasmid-mediated quinolone resistance gene, *qnrVC7*, in *Vibrio cholerae* of seafood origin. *International Journal of Antimicrobial Agents*. 2015;45(6):667-668. doi:[10.1016/j.ijantimicag.2015.02.002](https://doi.org/10.1016/j.ijantimicag.2015.02.002)
15. Pazhani GP, Chakraborty S, Fujihara K, et al. QRDR mutations, efflux system & antimicrobial resistance genes in enterotoxigenic *Escherichia coli* isolated from an outbreak of diarrhoea in Ahmedabad, India. *Indian Journal of Medical Research*. 2011;134:214-223.
16. Liu M, Wong MHY, Chen S. Mechanisms of fluoroquinolone resistance in *Vibrio parahaemolyticus*. *International Journal of Antimicrobial Agents*. 2013;42(2):187-188. doi:[10.1016/j.ijantimicag.2013.04.024](https://doi.org/10.1016/j.ijantimicag.2013.04.024)
17. Fang Y, Pan X, Lin L, et al. Molecular mechanisms of quinolone resistance in *Aeromonas hydrophila*. *Acta Microbiologica Sinica*. 2014;54:174-182.
18. Grabherr MG, Haas BJ, Yassour M, et al. Full-length transcriptome assembly from RNA-Seq data without a reference genome. *Nat Biotechnol*. 2011;29(7):644-652. doi:[10.1038/nbt.1883](https://doi.org/10.1038/nbt.1883). PMID:21572440

19. Conesa A, Götz S, García-Gómez JM, Terol J, Talón M, Robles M. Blast2GO: a universal tool for annotation, visualization and analysis in functional genomics research. *Bioinformatics*. 2005;21(18):3674-3676. doi:[10.1093/bioinformatics/bti610](https://doi.org/10.1093/bioinformatics/bti610)
20. Audic S, Claverie JM. The significance of digital gene expression profiles. *Genome Res*. 1997;7(10):986-995. doi:[10.1101/gr.7.10.986](https://doi.org/10.1101/gr.7.10.986)
21. Leng N, Dawson JA, Thomson JA, et al. EBSeq: an empirical Bayes hierarchical model for inference in RNA-seq experiments. *Bioinformatics*. 2013;29(8):1035-1043. doi:[10.1093/bioinformatics/btt087](https://doi.org/10.1093/bioinformatics/btt087). PMID:23428641
22. Willems E, Leyns L, Vandesompele J. Standardization of real-time PCR gene expression data from independent biological replicates. *Analytical Biochemistry*. 2008;379(1):127-129. doi:[10.1016/j.ab.2008.04.036](https://doi.org/10.1016/j.ab.2008.04.036)
23. Ogata H, Goto S, Sato K, Fujibuchi W, Bono H, Kanehisa M. KEGG: Kyoto Encyclopedia of genes and genomes. *Nucleic Acids Research*. 1999;28(1):29-34. doi:[10.1093/nar/28.1.27](https://doi.org/10.1093/nar/28.1.27). PMID:10592173
24. Li L, Feng S, Wang Y, Dai X, Yang Y, Zeng M. The transcriptome sequencing and analysis of *Salmonella typhimurium* ATCC13311 inducing drug-resistance strain. *Chinese Journal of Veterinary Science*. 2015;35:1088-1094.
25. Chang C, Stewart RC. The two-component system regulation of diverse signaling pathways in prokaryotes and eukaryotes. *Plant Physiology*. 1998;117(3):723-731. doi:[10.1104/pp.117.3.723](https://doi.org/10.1104/pp.117.3.723). PMID:9662515
26. Sperandio V, Torres AG, Kaper JB. Quorum sensing *Escherichia coli* regulators B and C (QseBC): a novel two-component regulatory system involved in the regulation of flagella and motility by quorum sensing in *E. coli*. *Molecular Microbiology*. 2002;43(3):809-821. doi:[10.1046/j.1365-2958.2002.02803.x](https://doi.org/10.1046/j.1365-2958.2002.02803.x)
27. Singh AK, Haldar R, Mandal D, Kundu M. Analysis of the topology of *Vibrio cholerae* NorM and identification of amino acid residues involved in norfloxacin resistance. *Antimicrob Agents Chemother*. 2006;50(11):3717-3723. doi:[10.1128/aac.00460-06](https://doi.org/10.1128/aac.00460-06). PMID:16954325
28. Bohnert JA, Karamian B, Nikaido H. Optimized Nile red efflux assay of AcrAB-TolC multidrug efflux system shows competition between substrates. *Antimicrob Agents Chemother*. 2010;54(9):3770-3775. doi:[10.1128/aac.00620-10](https://doi.org/10.1128/aac.00620-10). PMID:20606071
29. Long F, Rouquette-Loughlin C, Shafer WM, Yu EW. Functional cloning and characterization of the multidrug efflux pumps NorM from *Neisseria gonorrhoeae* and YdhE from *Escherichia coli*. *Antimicrob Agents Chemother*. 2008;52(9):3052-3060. doi:[10.1128/aac.00475-08](https://doi.org/10.1128/aac.00475-08). PMID:18591276
30. Chapman JS, Georgopapadakou NH. Routes of quinolone permeation in *Escherichia coli*. *Antimicrob Agents Chemother*. 1988;32(4):438-442. doi:[10.1128/aac.32.4.438](https://doi.org/10.1128/aac.32.4.438). PMID:3132091

## SUPPLEMENTARY MATERIALS

### **Supplementary Material for Xiaochen Tang and Yu Ding article - IJA**

Download: <https://ija.scholasticahq.com/article/84647-transcriptome-analysis-reveals-the-important-roles-of-a-two-component-system-flagellar-assembly-active-efflux-system-and-outer-membrane-proteins-in/attachment/173830.pdf>

---