

Original Research Articles

Transcriptome analysis of *Litopenaeus vannamei* during the early stage of limb regeneration process

Wucheng Yue^{1,2a}, Ruiming Yuan^{3b}, Donghuo Jang^{3,4}, Xiangzhao Guo⁵, Fuhua Li², Xueqiao Qian^{1c}

¹ Animal Husbandry and Fisheries Research Center of Guangdong Haid Group Co., Ltd. China, ² Institute of oceanology, Chinese academy of sciences. China, ³ Guangdong Haid Group Co., Ltd. China, ⁴ Hainan Haida Aquatic Seed Industry Development Co., Ltd. China, ⁵ Animal Husbandry and Fisheries Research Center of Guangdong Haid Group Co., Ltd. China 2Guangdong Haid Group Co., Ltd. China

Keywords: autotomy, regeneration, transcriptome

<https://doi.org/10.46989/001c.88325>

Israeli Journal of Aquaculture - Bamidgheh

Vol. 75, Issue 2, 2023

Regeneration is a process in which organisms regrow new tissues or organs at the injury site, which has attracted the attention of many scientists and nonscientists. However, the underlying molecular mechanisms of regeneration after autotomy are largely unknown. In this study, we conducted RNA-seq sequencing on regenerated limb bud tissues of *Litopenaeus vannamei* at 0 hours post autotomy (0 hpa), 12 hours post autotomy (12 hpa), and 24 hours post autotomy (24 hpa). A total of 2,192 differentially expressed genes related to energy metabolism, transcription and translation, and epidermis development were identified between 0 hpa and 12 hpa, such as triosephosphate isomerase A, triosephosphate isomerase B, and zinc finger protein 367 that is upregulated in 12 hpa. Between 12 hpa and 24 hpa, 1,447 differentially expressed genes were identified that were related to cuticle development and energy metabolism, such as cuticle protein 6, which is upregulated in 24 hpa, and triosephosphate isomerase is downregulated in 24 hpa. The results indicated that energy metabolism, transcription and translation, epidermal formation, and chitin metabolism processes are involved during the early stage of limb regeneration. This study provides basic knowledge for investigating the molecular mechanisms associated with limb regeneration in crustaceans at the early regeneration stage.

INTRODUCTION

Regeneration is when organisms replace or restore lost or amputated body parts. The regeneration capacities are different in various organisms, based mainly on sex, age, and different development stages,^{1,2} and invertebrate species show overall superior regeneration ability compared to vertebrates.³ For example, starfish can regrow a new individual from a small body fragment,⁴ lizard can regenerate a new tail after its tail self-cut.⁵ Regeneration enables animals to recover well despite substantial injuries or autotomy. It is an important biological process for some animals to survive after injuries.

Regeneration involves a series of processes. The early stage is wound healing, which includes the process of the wound epidermis formation and recovery of body fluid. Then, the blastema is formed, that is a proliferative zone of cells consisting of mesenchymal progenitor cells. Finally, a redevelopment phase follows, when tissue grows again.⁶

Early stages of regeneration are critical for autotomy signal transduction and determine whether an amputated limb can regenerate successfully. There are many studies focused on the role of single molecules in the early stage of regeneration. In *Xenopus laevis*, Wnt-3a has played an role in blastema formation in the apical epithelium of regenerating limb buds.⁷ *IGF-I* is an important gene for controlling the regeneration of retinal ganglion cells during the early stage of regeneration in goldfish.⁸ *Fgf* is required to form blastema at the early stage of fin regeneration and then activate the regenerative cell proliferation in zebrafish.⁹ However, the molecular mechanism of animal regeneration during the early stage at transcriptome level is still incomplete.

Regeneration is a typical biological character in crustaceans; however, previous studies have mainly focused on the morphology of newly formed limbs and the effect of autotomy on regeneration phenotype in crustaceans. In *Parhyale hawaiiensis*, the regenerated limb appears to be a perfect replica of the original limb before amputation.¹⁰

a a These authors contributed equally to this study

b b These authors contributed equally to this study

c c Corresponding author. Xueqiao Qian; email: xwchenle@163.com

The size of the newly regenerated limb is the same as that of the normal limb after the second molt in *Eriocheir sinensis*.¹¹ In *Alpheus angulosus*, the regenerated claw differs from mature claws in the first molt, while the most of differences disappeared in the second molt.¹² No matter whether partial or full limb autotomy, it had no significant impact on the survival rate in *Scylla serrata*.² However, the molecular mechanism of limb regeneration in crustaceans is still not fully understood currently.

Litopenaeus vannamei is widely distributed in the world and is also the most economically valuable cultured crustacean.¹³ During the breeding process of *L. vannamei*, regeneration occurs frequently. However, the severed limbs could grow into new limb buds and eventually form new limbs. In this study, a comparative transcriptomics study of limb regeneration of *L. vannamei* was carried out to identify potential genes associated with limb regeneration at the early stage (less than 24 hours). This study aimed to help us understand the underlying molecular mechanism of limb regeneration of *L. vannamei* at early stage.

MATERIALS AND METHODS

EXPERIMENTAL ANIMALS AND SAMPLING

All the male *L. vannamei* individuals (~10g) in this study were collected from the Jinshan breeding station in Shanghai. First, they were acclimatized in the same tank for two days and then were physically stimulated on the second limb on the right side and autotomy took place at the coxa. Then, the limb bud tissues at the coxa from the wound site were sampled separately at 12 hours (12hpa) and 24 hours post autotomy (24hpa). The bases of coxae were collected from the second limb on the right side at 0-hour post autotomy (0hpa) for the control group. For each group, four individuals were sampled as biological replicates. Finally, the samples were immediately placed in liquid nitrogen and stored at -80°C in a refrigerator. In this study, no rare or protected species were used.

RNA EXTRACTION AND RNA SEQUENCING

Total RNA was extracted from the collected tissues of each individual by using RNAios plus (Takara, Dalian) according to the manufacturer's instructions. RNA integrity and quantity were examined using agarose gel electrophoresis and an Agilent 2100 Bioanalyzer (Agilent, Shanghai, China). A total of 5 µg RNA with RNA integrity number (RIN) above 8.0 was used for RNA-seq library construction using the Illumina Truseq™ RNA sample Prep Kit (Illumina, USA). Then, indexed libraries were sequenced on Illumina HiSeq™4000, with 150 bp paired-end reads produced.

IDENTIFICATION OF DIFFERENTIALLY EXPRESSED GENES

The raw sequencing reads were first trimmed to remove low-quality reads using Trimmomatic software. Clean reads were mapped against the assembled *L. vannamei* genome (NCBI accession number: GCF_003789085.1) using

hisat2-2.0 software. FPKM (fragments per kilobase per transcript per million mapped reads) value was calculated with the RSEM software for each individual to identify the expression levels. The differentially expressed genes (DEGs) were identified with a fold change > 2 and with statistical significance ($p < 0.05$) by DESeq2 software. Cluster analysis was conducted using the hierarchical cluster method based on the euclidean distance using the heatmap module in R.

FUNCTIONAL ANNOTATION

Functional annotation was performed by searching NR (NCBI nonredundant protein sequences), Swiss-Prot (A manually annotated and reviewed protein sequence database), GO (Gene Ontology), COG (Cluster of Orthologous Groups of proteins), and KEGG (Kyoto Encyclopedia of Genes and Genomes) databases using DIAMOND software.

WEIGHTED CORRELATION NETWORK ANALYSIS (WGCNA) ANALYSIS AND GO ANALYSIS

Based on the differentially expressed genes, a WGCNA co-expression network was constructed. The weighted signed network was computed based on a fit to scale-free topology, with a threshold soft power of 6. A topological overlap dendrogram was used to define modules with a minimum module size of 30 genes and the deepSplit parameter set to 2. Module and trait associations were determined by calculating the correlation coefficient between module eigengene (ME) and traits (such as days after limb autotomy). GO enrichment analysis was conducted through Goatools software with $p < 0.05$. KEGG enrichment analysis were conducted on the DEGs by DAVID functional annotation tools 6.8. The p-values were adjusted using Fisher exact test. And the categories with adjusted p-value < 0.05 were considered enriched.

RESULTS

ANALYSIS OF SEQUENCING QUALITY

Most of the genes with a quantitative value of more than 3.5 are close to saturation when the rate of map sequencing reads is 40% (Figure 1A). This means that the overall quality of saturation is high, and the sequencing amount of transcriptome can cover most of the expressed genes in this study. The curve in Figure 1B reflects that the sequence obtained by sequencing is evenly distributed on the gene, and the sequencing is not biased. The results prove that our transcriptome data are available.

DIFFERENTIALLY EXPRESSED GENES

A total of 2,192 DEGs were identified between 0 hpa and 12 hpa ($P < 0.05$, fold change >2), with 1,237 up-regulated and 955 down-regulated genes in the 12 hpa group. And a total of 1,447 DEGs were identified between 12 hpa and 24 hpa groups, with 368 up-regulated and 1,079 down-regulated genes in 12 hpa group (Figure 2).




Figure 1. The figure of sequencing quality assessment.

A. Saturation curve of sequencing. B. Coverage of sequencing.

Compared to 0 hpa group, the expression of genes involved in glucose metabolism upregulated significantly in the 12 hpa group, such as triosephosphate isomerase B-like (*tpi1b*), triosephosphate isomerase A-like (*tpi1a*), facilitated trehalose transporter Tret1-like (*Tret1*), sodium/glucose co-transporter 5-like (*SLC5A1*). The expression of genes related to transcription and translation also upregulated significantly in the 12 hpa group, such as elongation factor 2-like (*eEF2*), zinc finger protein 367-like (*znf367*), friend leukemia integration 1 transcription factor-like (*Fli1*). Besides, the expression of genes related to epidermal formation was also significantly upregulated in the 12 hpa group, such as innexin *inx3*-like (*Inx3*) and innexin *inx2*-like (*Inx2*). Conversely, the expression of genes related to cuticle and crustacyanin downregulated in the 12 hpa group, such as larval cuticle protein 16/17-like, cuticle protein CP1499-like, crustacyanin-A2 subunit-like, and crustacyanin-C1 subunit-like (Table 1).

Compared to the 12 hpa group, the expression of genes related to muscle is upregulated, and related to glucose metabolism are downregulated in the 24 hpa group, such as titin (*TTN*) and triosephosphate isomerase-like (*tpi1a*). Interestingly, from 12 hours to 24 hours after autotomy, some cuticle-related genes are highly expressed, and some are low expressed. For instance, cuticle protein 6 and cuticle protein CP575-like are upregulated, and pupal cuticle protein 20-like and cuticle protein 8-like are downregulated in the 24 hpa group (Table 2).

WGCNA AND FUNCTIONAL ENRICHMENT ANALYSIS

WGCNA analysis indicated that a total of 55 differentially expressed genes are significantly related to the 12 hpa group (grey module), and a total of 48 differentially expressed genes are significantly related to the 24 hpa group (pink module) (Figure 3A). Gene ontology enrichment analysis was used to classify these differentially expressed




Figure 2. The profile of differentially expressed genes during the early stage of limb regeneration of *L.vannamei*.

A. Volcano plot of differentially expressed genes between 0 hpa and 12 hpa and between 12 hpa and 24 hpa. The green dots indicate that the gene is downregulated. The red dots indicate that the gene is upregulated. The gray dots indicate that there was no significant difference. B. Heatmap plot of differentially expressed genes during the early stage of the limb regeneration process. The blue color indicates low expression, and the red color indicates high expression.

genes. The 55 differentially expressed genes in the grey module were mainly enriched in serine hydrolase activity, carbohydrate metabolic process, and generation of precursor metabolites and energy. And the 48 differentially expressed genes in the pink module were mainly enriched in the plasma membrane region, regulation of ion transmembrane transport, and voltage-gated channel activity (Figure 3B, C).

KEGG enrichment analysis was also conducted on the DEGs and the top 10 significantly enriched KEGG pathways were presented. The differentially expressed genes between 0 hpa and 12 hpa group mainly enriched in lipid and atherosclerosis, inositol phosphate metabolism, glycolysis/gluconeogenesis, fructose and mannose metabolism, ribosome biogenesis in eukaryotes and pathogenic infection KEGG pathways. The differentially expressed genes between the 12 hpa group and 24 hpa group mainly enriched in glycosaminoglycan biosynthesis, complement and coagulation cascades, and renin-angiotensin system pathways (Figure 4).

DISCUSSION

The capacity to regenerate new limb after autotomy is a surprising thing and most crustaceans have this ability, such as *Uca pugilator*,¹⁴ *Faxonius virilis*,¹⁵ *Eriocheir sinensis*.¹⁶ In this study, the transcriptome of *L. vannamei* was sequenced at the early stage of limb regeneration (0 hours, 12, hours and 24 hours post autotomy), and the differentially expressed genes were analyzed. The results indicated that regeneration after autotomy involves energy metabolism, epithelial formation, transcription and translation, and cuticle metabolism processes in *L. vannamei*.

Compared with 0 hpa group, the most significant changes in gene expression at 12 hpa are the genes related to energy metabolism, such as triosephosphate isomerase A, triosephosphate isomerase B, sodium/glucose cotransporter 5, and so on at 12 hpa (Table 1). Regeneration of the autotomized chelipeds imposes an additional energy demand which is called regeneration load in crustaceans.¹⁷ Animals need to acquire sufficient energy to regenerate

Table 1. The genes with significant expression change at 0 hour (0 hpa) and 12 hours post autotomy group (12 hpa).

Gene ID	Gene	Uniprot	Fold Change (12 hpa/0 hpa)	Function
LOC113801022	triosephosphate isomerase B-like	Q90XG0	253.677	Glycolysis and Gluconeogenesis.
LOC113802426	triosephosphate isomerase A-like	Q1MTI4	251.024	Glycolysis and Gluconeogenesis.
LOC113825018	facilitated trehalose transporter Tret1-like	A9ZSY2	73.29	Regulating trehalose levels in the hemolymph.
LOC113828571	sodium/glucose cotransporter 5-like	P53791	18.004	Transport of dietary monosaccharides.
LOC113811541	elongation factor 2-like	P13060	411.348	Translation elongation.
LOC113819444	H/ACA ribonucleoprotein complex subunit 1-like	Q7KVQ0	376.358	Required for ribosome biogenesis.
LOC113823896	zinc finger protein 367-like	Q567C6	53.834	Transcriptional activator.
LOC113806657	transcriptional regulator Erg-like	Q01543	25.413	Sequence-specific transcriptional activator.
LOC113802698	Friend leukemia integration 1 transcription factor-like	P26323	24.337	Transcriptional activator.
LOC113800864	innexin inx3-like	Q9VAS7	18.928	Essential for proper epithelial development of the epidermis.
LOC113800859	innexin inx2-like	Q9V427	18.421	Required for epithelial morphogenesis.
LOC113804810	ETS homologous factor-like	O70273	12.655	Regulating epithelial cell differentiation and proliferation.
LOC113824715	crustacyanin-A1 subunit-like	A0A8J5J9V4	0.057	Provides the blue coloration to the carapace.
LOC113829675	crustacyanin-A2 subunit-like	P80007	0.01	Provides the blue coloration to the carapace.
LOC113829725	crustacyanin-C1 subunit-like	P80029	0.015	Provides the blue coloration to the carapace.
LOC113821825	cuticle protein CP575	P81589	0.045	Cuticle
LOC113829880	cuticle protein 6	P82119	0.036	Cuticle
LOC113804370	cuticle protein CP1499-like	P81584	0.033	Cuticle

*Just listed the genes with significant expression changes at the top.

Table 2. The genes with significant expression change in 12 hours post autotomy group (12 hpa) and 24 hours post autotomy group (24 hpa).

Gene ID	Gene	Uniprot	Fold Change (24 hpa/12 hpa)	Function
LOC113821794	cuticle protein CP575	P81589	150.7	Cuticle
LOC113827056	cuticle protein 6	P82119	110.129	Cuticle
LOC113816627	pupal cuticle protein 20-like	O02388	0.019	Cuticle
LOC113825526	cuticle protein 8-like	Q9V7U0	0.011	Cuticle
LOC113801304	titin	Q8WZ42	79.175	Key component of muscles.
LOC113820944	tropomyosin-like	Q7M3Y8	18.03	Regulation of contraction
LOC113801022	triosephosphate isomerase B-like	Q90XG0	0.122	Glycolysis and gluconeogenesis
LOC113800990	triosephosphate isomerase-like	Q1MTI4	0.063	Glycolysis and gluconeogenesis

*Just listed the genes with significant expression changes at the top.

the lost limb and will allocate energy after autotomy.¹⁸ These results can explain why the expression of genes related to energy metabolism upregulates rapidly at 12 hpa.

In addition, the expression of genes related to transcription and translation also upregulated sharply at 12 hpa, such as zinc finger protein 367-like and transcriptional regulator




Figure 3. WGCNA and GO enrichment analysis.

A. Relationships of consensus module eigengenes and limb regeneration stage identified by WGCNA analysis. Highest correlation coefficients and small p-values indicate that the co-expressed genes in the module were strongly associated with the regeneration stages. B. GO enrichment analysis for genes strongly associated with 12 hpa by WGCNA analysis. C. GO enrichment analysis for genes strongly associated with 24 hpa by WGCNA analysis.

Erg (Table 1). The process of regeneration requires a high amount of transcriptional and translational regulation.¹⁹ Transcriptional activity increases significantly in Echinoderms during the early stages of regeneration,²⁰ and elongation factor is upregulated during arm regeneration in *Amphiura filiformis*.¹⁹ The expression of transcription and translation related genes upregulated after autotomy in *L. vannamei* were reasonable. The results in this study indicated that the expression of genes involved in energy production and transcription and translation processes changed significantly during the early stages of regeneration is consistent with the study in *A. filiformis*.²¹ Energy production and transcription and translation processes are essential in restoring the volume of coelomic fluid and the number of coelomocytes after autotomy that is the first response of regeneration.²² Besides, the genes related to epidermal development are also upregulated at 12 hpa, such

as innexin *inx2*-like, innexin *inx3*-like and ETS homologous factor-like. Innexin 2 is upregulated during the limb regeneration process and likely facilitates the early stage of the regeneration process in *E. sinensis*.²³ Innexin mediates instructive signaling and participated in the regeneration process of *D. japonica*.^{24,25} Innexin may play a crucial role in the regeneration process within 12 hpa in *L. vannamei*. In sum, the genes that are related to energy metabolism, transcription and translation, and epithelial development are activated in the regeneration process after the autotomy of *L. vannamei*.

From 12 hours to 24 hours after autotomy, the expression of genes that were related to the cuticle and energy metabolism changed significantly, such as cuticle protein 6 and triosephosphate isomerase (Table 2). The expression of cuticle protein 6 and cuticle protein CP575 upregulated, and cuticle protein downregulated during 12 hours to 24




Figure 4. KEGG enrichment analysis of differentially expressed genes with top 10 enriched pathways presented.

A. KEGG enrichment analysis of upregulated genes at 12 hpa compared to 0 hpa. B. KEGG enrichment analysis of downregulated genes at 12 hpa compared to 0 hpa. C. KEGG enrichment analysis of upregulated genes at 24 hpa compared to 12 hpa. D. KEGG enrichment analysis of downregulated genes at 24 hpa compared to 12 hpa.

hours after autotomy, while the expression of cuticle protein 6 and cuticle protein CP575 downregulated at 12 hpa compared to 0 hpa group. In *Portunus trituberculatus*, cuticle proteins were up regulated during limb regeneration, suggesting the cuticle proteins might promote regeneration processes, which is consistent with our research.²⁶ Conversely, the expression of some cuticle proteins downregulated in regeneration processes. The reason may be different cuticle protein genes performed different functions. Researchers have found that cuticle protein genes have different spatial and temporal expressions and may perform different functions in *Cherax quadricarinatus* and *Locusta migratoria*.^{27,28} Thus, it needs further experimental evidence to confirm the function of cuticle protein in the regeneration process.

In conclusion, regeneration after autotomy is a complex and dynamic process. At the early stage (less than 24 hours), limb regeneration involves energy metabolism, transcription and translation, epidermal formation, and cuticle metabolism process in *L. vannamei*.

COMPETING INTERESTS

The authors have declared no competing interests.

ACKNOWLEDGMENTS

This work was supported by Guangdong International Science & Technology Cooperation Program 2020 (2020A0505100026).

AUTHORS' CONTRIBUTION

Conceptualization: Wucheng Yue (Equal), Fuhua Li (Equal). Formal Analysis: Wucheng Yue (Equal), Ruiming Yuan (Equal). Investigation: Wucheng Yue (Equal), Ruiming Yuan (Equal). Writing – original draft: Wucheng Yue (Equal), Ruiming Yuan (Equal). Resources: Donghuo Jang (Equal), Xiangzhao Guo (Equal). Writing – review & editing: Xueqiao Qian (Lead). Funding acquisition: Xueqiao Qian (Lead). Supervision: Xueqiao Qian (Lead).

Submitted: June 14, 2023 CST. Accepted: September 26, 2023
CST. Published: October 04, 2023 CST.



This is an open-access article distributed under the terms of the Creative Commons Attribution 4.0 International License (CCBY-4.0). View this license's legal deed at <http://creativecommons.org/licenses/by/4.0> and legal code at <http://creativecommons.org/licenses/by/4.0/legalcode> for more information.

REFERENCES

1. Jeffery WR. Distal regeneration involves the age dependent activity of branchial sac stem cells in the ascidian *Ciona intestinalis*. *Regeneration*. 2014;2(1):1-18. doi:[10.1002/reg2.26](https://doi.org/10.1002/reg2.26). PMID:25893097
2. Quintio ET, Estepa FDP. Survival and growth of mud crab, *Scylla serrata*, juveniles subjected to removal or trimming of chelipeds. *Aquaculture*. 2011;318(1-2):229-234. doi:[10.1016/j.aquaculture.2011.05.034](https://doi.org/10.1016/j.aquaculture.2011.05.034)
3. Slack JMW. Animal regeneration: Ancestral character or evolutionary novelty? *EMBO Reports*. 2017;18(9):1497-1508. doi:[10.15252/embr.201643795](https://doi.org/10.15252/embr.201643795). PMID:28747491
4. Byrne M. The link between autotomy and CNS regeneration: Echinoderms as non-model species for regenerative biology. *BioEssays*. 2020;42:1900219. doi:[10.1002/bies.201900219](https://doi.org/10.1002/bies.201900219)
5. Alibardi L. Ultrastructural analysis of early regenerating lizard tail suggests that a process of dedifferentiation is involved in the formation of the regenerative blastema. *J Morphol*. 2018;279(8):1171-1184. doi:[10.1002/jmor.20838](https://doi.org/10.1002/jmor.20838)
6. Han M, Yang X, Taylor G, Burdsal CA, Anderson RA, Muneoka K. Limb regeneration in higher vertebrates: Developing a roadmap. *Anat Rec B New Anat*. 2005;287B(1):14-24. doi:[10.1002/ar.b.20082](https://doi.org/10.1002/ar.b.20082)
7. Yokoyama H, Ogino H, Stoick-Cooper CL, Grainger RM, Moon RT. Wnt/ β -catenin signaling has an essential role in the initiation of limb regeneration. *Dev Biol*. 2007;306(1):170-178. doi:[10.1016/j.ydbio.2007.03.014](https://doi.org/10.1016/j.ydbio.2007.03.014). PMID:17442299
8. Koriyama Y, Homma K, Sugitani K, et al. Upregulation of igf-I in the goldfish retinal ganglion cells during the early stage of optic nerve regeneration. *Neurochem Int*. 2007;50(5):749-756. doi:[10.1016/j.neuint.2007.01.012](https://doi.org/10.1016/j.neuint.2007.01.012)
9. Shibata E, Yokota Y, Horita N, et al. Fgf signalling controls diverse aspects of fin regeneration. *Development*. 2016;143(16):2920-2929. doi:[10.1242/dev.140699](https://doi.org/10.1242/dev.140699)
10. Almazán A, Çevrim Ç, Musser JM, Averof M, Paris M. Crustacean leg regeneration restores complex microanatomy and cell diversity. *Sci Adv*. 2022;8(34):eabn9823. doi:[10.1126/sciadv.abn9823](https://doi.org/10.1126/sciadv.abn9823). PMID:36001670
11. He J, Wu X, Cheng Y. Effects of limb autotomy on growth, feeding and regeneration in the juvenile *Eriocheir sinensis*. *Aquaculture*. 2016;457:79-84. doi:[10.1016/j.aquaculture.2016.02.004](https://doi.org/10.1016/j.aquaculture.2016.02.004)
12. Pereira A, Tracey E, Cooney PC, Korey CA, Hughes M. Post-autotomy claw regrowth and functional recovery in the snapping shrimp *Alpheus angulosus*. *Mar Freshw Behav Phy*. 2014;47(3):147-159. doi:[10.1080/10236244.2014.928460](https://doi.org/10.1080/10236244.2014.928460)
13. Wang Z, Wu Q, Liao G, Fan L. New insights into the regulation mechanism of *Litopenaeus vannamei* hepatopancreas after lipopolysaccharide challenge using transcriptome analyses. *Fish Shellfish Immun*. 2022;128:466-473. doi:[10.1016/j.fsi.2022.08.014](https://doi.org/10.1016/j.fsi.2022.08.014)
14. McLain DK, Pratt AE. Body and claw size at autotomy affect the morphology of regenerated claws of the sand fiddler crab, *Uca pugnator*. *J Crustacean Biol*. 2011;31(1):1-8. doi:[10.1651/10-3298.1](https://doi.org/10.1651/10-3298.1)
15. Graham ZA, Vargas C, Angilletta MJ Jr, Palaoro AV. Regenerated claws of the virile crayfish *Faxonius virilis* (Hagen, 1870) (Decapoda: Astacidea: Cambaridae) generate weaker pinching forces compared to original claws. *Journal of Crustacean Biology*. 2021;41(3):ruab036. doi:[10.1093/jcbiol/ruab036](https://doi.org/10.1093/jcbiol/ruab036)
16. Li J, Tian Y, Li X, Zuo J, Zhao R, Sun J. Insulin-like signaling promotes limb regeneration in the Chinese mitten crab (*Eriocheir sinensis*). *Fish Shellfish Immun*. 2022;122:268-275. doi:[10.1016/j.fsi.2022.02.007](https://doi.org/10.1016/j.fsi.2022.02.007)
17. Mariappan P, Balasundaram C, Schmitz B. Decapod crustacean chelipeds: An overview. *J Biosci*. 2000;25(3):301-313. doi:[10.1007/bf02703939](https://doi.org/10.1007/bf02703939)
18. Devi S a A, Rita G. Effect of multiple limb autotomy on oogenesis and somatic growth in *paratelphusa hydromous*. *Trop fres biol*. 2000;9:43-56.
19. Purushothaman S, Saxena S, Meghah V, et al. Transcriptomic and proteomic analyses of *Amphiura filiformis* arm tissue-undergoing regeneration. *J Proteomics*. 2015;112:113-124. doi:[10.1016/j.jprot.2014.08.011](https://doi.org/10.1016/j.jprot.2014.08.011)
20. Quispe Parra DJ. *Differential Gene Expression during Early Stages of Regeneration in Holothuria Glaberrima*. University of Puerto. Rio Piedras; 2021.

21. Burns G, Ortega-Martinez O, Thorndyke MC, Peck LS, Dupont S, Clark MS. Dynamic gene expression profiles during arm regeneration in the brittle star *amphiura filiformis*. *J Exp Mar Biol Ecol*. 2011;407(2):315-322. doi:[10.1016/j.jembe.2011.06.032](https://doi.org/10.1016/j.jembe.2011.06.032)
22. Dolmatov IY. Molecular aspects of regeneration mechanisms in holothurians. *Genes (Basel)*. 2021;12(2):250. doi:[10.3390/genes12020250](https://doi.org/10.3390/genes12020250). PMID:33578707
23. Wang J, Chen X, Hou X, et al. “Omics” data unveil early molecular response underlying limb regeneration in the Chinese mitten crab, *Eriocheir sinensis*. *Sci Adv*. 2022;8(37):eabl4642. doi:[10.1126/sciadv.abl4642](https://doi.org/10.1126/sciadv.abl4642). PMID:36112682
24. Nogi T, Levin M. Characterization of innexin gene expression and functional roles of gap-junctional communication in planarian regeneration. *Dev Biol*. 2005;287(2):314-335. doi:[10.1016/j.ydbio.2005.09.002](https://doi.org/10.1016/j.ydbio.2005.09.002)
25. Oviedo NJ, Morokuma J, Walentek P, et al. Long-range neural and gap junction protein-mediated cues control polarity during planarian regeneration. *Dev Biol*. 2010;339:188-199. doi:[10.1016/j.ydbio.2010.03.018](https://doi.org/10.1016/j.ydbio.2010.03.018)
26. Fu Y, Zhu F, Liu L, et al. Itraq-based proteomic analysis identifies proteins involved in limb regeneration of swimming crab *Portunus trituberculatus*. *Comp Biochem Phys D*. 2018;26:10-19. doi:[10.1016/j.cbd.2018.02.003](https://doi.org/10.1016/j.cbd.2018.02.003)
27. Zhao X, Gou X, Qin Z, et al. Identification and expression of cuticular protein genes based on locusta migratoria transcriptome. *Sci Rep*. 2017;7(1):45462. doi:[10.1038/srep45462](https://doi.org/10.1038/srep45462). PMID:28368027
28. Zhou S, Jiang K, Liu M, Wang B, Wang L. Exploration of the functional properties of red cuticle patch of redclaw crayfish (*Cherax quadricarinatus*) by histological and transcriptomic analysis. *Aquaculture*. 2022;561:738624. doi:[10.1016/j.aquaculture.2022.738624](https://doi.org/10.1016/j.aquaculture.2022.738624)