

Original Research Articles

Application of fuzzy model for the shelf-life determination of whole sea bass (*Dicentrarchus labrax*) under refrigerated temperatures

İsmail Y. Genç¹0^a

¹ Fishing and Processing Technology Department, Eğirdir Fisheries Faculty, Isparta University of Applied Sciences Keywords: Fuzzy logic, shelf life, modeling, seafood, quality, seabass https://doi.org/10.46989/001c.88490

Israeli Journal of Aquaculture - Bamidgeh

Vol. 75, Issue 2, 2023

In this study, microbiological deterioration of whole sea bass stored at refrigerator temperatures (4°C) was determined through fuzzy modeling. Total Mesophilic Aerobic Bacteria (TMAB), Total Psychrophilic Aerobic Bacteria (TPAB), Lactic Acid Bacteria (LAB), Pseudomonas sp., and Enterobacteriaceae counts were observed during 12 days of storage. Quality Index Method (QIM) was used for sensory analysis of the samples and shelf life was determined according to sensory deterioration. The growth of the microorganisms was statistically significant (p<0.05) at the end of the storage period. Additionally, significant differences (p<0.05) were observed for QI scores after the 8th day of the storage in whole sea bass stored under refrigerated conditions. A total of 12 rules were determined in the fuzzy logic application. Mamdani was used as the fuzzy type for fuzzy modeling, and centroid was used as the defuzzification method. Fuzzy membership values were computed using triangular membership functions. According to the research results, the shelf life of sea bass fish was determined as 8 days from the time of 12 days storage. As a result, it was observed that the sensory score could be determined by using Fuzzy logic (R²=0.96) according to microbiological parameters in sea bass stored in refrigerator conditions. In this context, it was concluded that Fuzzy logic could be used in shelf-life estimation in seafood products.

INTRODUCTION

Quality in seafood products is described as changes influenced by product characteristics and environmental conditions.¹ Moreover, the freshness of raw fish has been reported as the most important criterion for consumer acceptability and as a parameter that can be determined by organoleptic evaluation.² Besides, microbiological, chemical, and sensory parameters are used to determine the quality of seafood, and gutted and ungutted status also affects the quality changes of seafood products during storage.³ Temperature is a significant factor during seafood storage, especially in slowing or preventing microbial growth and maintaining sensory quality.⁴

Microbial spoilage of seafood is caused by specific spoilage organisms (SSOs). Due to the nutrient content, pH, and weak connective tissues in seafood, SSOs grow very intensely during storage, and these spoilage-causing microorganisms lead to off-odors and quality losses. Their growth is related to quality. The microorganisms that are reported to be responsible for the spoilage of seafood belong to the genera *Shewanella*, *Pseudomonas*, *Photobacterium*, *Acinetobacter*, *Moraxella*, and *Psychrobacter*.⁵⁻⁷ Sen-

sory evaluation of seafood is the assessment by assessors of the changes that occur in seafood during storage and that can be perceived by the senses, such as sight, smell, taste, or touch. In this context, sensory evaluation provides information about the sensory quality of the product and also allows evaluation of the storage time.⁸ The QIM is a prominent way of determining the freshness of seafood based on a limited set of sensory characteristics. These features are graded from 0 to 3, with the aggregate determining the QI. As a result, the linear connection between QI and storage period and the shelf-life may be estimated. Furthermore, QIM is considered specific to each seafood species.⁹

In addition to microbiological, chemical, and sensory methods traditionally used in determining the quality of seafood products, mathematical models such as predictive microbiology¹⁰ and artificial neural networks¹¹ can also be used in determining the quality of seafood products. However, a limited number of studies determine the quality changes in seafood using fuzzy logic.^{12,13} The fuzzy logic represents the degree of incomprehensibility in the human mind by connecting it to a real number and dealing with uncertain events quantitatively.¹⁴ In this context, ease of use in biological processes such as microbiological and sen-

sory changes is considered one of the major advantages. In this framework, this study aims to develop a Fuzzy model by using microbiological and sensory changes in whole sea bass stored under refrigerator conditions as input and output, respectively, and to determine the shelf life according to the developed Fuzzy model.

MATERIALS AND METHODS

MATERIALS

This study used 24 sea bass with a total weight of 327.29 ± 23.26 g. The samples were obtained from a commercial establishment and delivered in ice-protected styrofoam boxes to the laboratory. The samples were stored aerobically under refrigerator conditions (+4°C) for 12 days and were covered with plastic bags to prevent drying of the skins due to the refrigerator conditions. Microbiological and sensory analyses were performed on days 0,2,5,8, and 12 of storage.

METHODS

MICROBIOLOGICAL ASSESSMENT

Microbiological analyses were periodically performed on whole sea bass stored under refrigerated conditions. On each sampling day, 25 g samples were taken aseptically and 10-fold dilutions were prepared. TMAB and TPAB were incubated on Plate Count Agar (PCA, Merck) at 30 and 4°C for 48 and 240 hours, respectively. Lactic Acid Bacteria (LAB) were enumerated on De Man Rogosa and Sharp (MRS) agar (Merck), *Pseudomonas sp.* were counted on Cephaloridin-Fucidin-Cetrimide (CFC, Merck) agar with CFC supplement and finally Enterobacteriaceae were determined on Violate Red Bile Glucose (VRBG, Merck) agar. LAB, *Pseudomonas sp.*, and members of the Enterobacteriaceae family were incubated at 25°C for 120 h, 25°C for 48 h, and 25°C for 36-48 h, respectively. Bacterial counts were expressed as log cfu/ g.¹⁵

SENSORY ANALYSIS

The sensory analysis of the whole sea bass, which were stored under refrigerator conditions, was carried out according to the Quality Index Method (QIM) proposed by Knowles et al.¹⁶ In this context, skin brightness, hemorrhage on skin and fins, hardness, eye structure, odor and color of gills, abdominal color and vent condition parameters were evaluated by five trained assessors. The assessors gave a score between 0 and 2 or 3 for each demerit point and continued up to 22 points as a total QI score.

FUZZY MODELING

Fuzzy modeling is a method proposed by Zadeh.¹⁷ In this context, it consists of 3 main steps: fuzzification, fuzzy process, and defuzzification. In this study, fuzzy modeling specified by Zadeh¹⁷ was applied to sea bass according to the method specified by Zare and Ghazali.¹² In this regard, inputs and outputs were introduced to the fuzzy system

in the fuzzification step. The TMAB, TPAB, Pseudomonas sp., LAB, and Enterobacteriaceae numbers were used as input. Microorganism counts were introduced to the system as low, medium, and high. In the fuzzy processing step, the IF...THEN, rules necessary for the establishment of the model were generated. A total of 12 rules were formed for whole sea bass stored under refrigerated conditions. QI scores, which are output parameters, were matched with the input parameters within the framework of the rules. One of the rules is as follows;

IF TMAB is low, TPAB is low, LAB is low, Pseudomonas sp. is medium, and Enterobacteriaceae is low, THEN QI score is low.

Finally, in the defuzzification step, an output value was calculated by fuzzy logic according to the rules generated in the fuzzy processing part, where each rule corresponds to an output parameter (QI) corresponding to a different scenario. To calculate according to the input parameters and the rules defined, the output parameters must also have a category as low, medium, and high. In this section, the centroid method is used for defuzzification.

STATISTICAL ANALYSIS

In this study, the analysis of variance was calculated using the ANOVA test. Tukey's multiple analysis test was used for significant differences between means. The results are given as mean ± standard deviation. SPSS (SPSS, USA) package program was used for statistical analyses. MATLAB R2014a (The MathWorks, USA) was used for fuzzy modeling.

RESULTS

MICROBIOLOGICAL ASSESSMENT

The Microbiological growth of whole sea bass stored aerobically under refrigerated conditions is shown in Fig 1. In this context, TMAB counts were determined on the first day of storage to be $3.16\pm0.12 \log \text{cfu/g}$. Statistically significant increases were found at the beginning and end of storage (p<0.05) and TMAB counts were determined to be 6.53 ± 0.76 log cfu/g on the last day of storage. No statistical difference was observed on the 5th, 8th and 12th day of storage (p>0.05). The increase in TPAB counts was similar to that of TMAB (Figure 1). Accordingly, TPAB counts were $2.50\pm0.70 \log \text{cfu/g}$ on the first day of storage and significantly (p<0.05) increased to $7.31\pm0.08 \log \text{cfu/g}$ at the end of storage. No statistically significant difference was observed on the 8th and 12th day of storage (p>0.05).

Pseudomonas sp. numbers were determined to be the highest at the beginning and end of storage compared to other bacterial groups. It was observed that *Pseudomonas sp.* numbers, initially determined as 3.32 ± 0.25 log cfu/g at the beginning of storage, increased statistically significantly (p<0.05) and reached 7.60±0.31 log cfu/g at the end of storage. *Pseudomonas sp.* showed a statistically significant increase during the storage period and no significant differences were determined between the 8th and 12th days (p>0.05). Lactic acid bacteria and members of the Enter-



Figure 1. Microbiological assessment of whole seabass stored under refrigerated temperatures.

obacteriaceae family showed slower growth than other bacterial groups in aerobically stored sea bass under refrigerator conditions. Therefore, LAB and Enterobacteriaceae counts on the first day of storage were 2.25 ± 0.49 and $1.65\pm0.49 \log$ cfu/g, respectively. While no statistical difference was observed for LAB counts on the 5th and 8th day of storage (p>0.05), a significant increase (p<0.05) was observed on the 12th day, the last day of storage, reaching $4.58\pm0.04 \log$ cfu/g. No statistical differences were observed in the members of the Enterobacteriaceae family on the 5th, 8th and 12th days of storage and their number was determined to be $4.27\pm0.14 \log$ cfu/g on the last day of storage. However, significant increases were observed compared to the initial day of storage (p<0.05).

SENSORY CHANGES

Sensory changes in sea bass aerobically stored under refrigerator conditions are shown in Figure 2. Accordingly, the samples evaluated in terms of skin brightness, hemorrhage on skin and fins, hardness, eye structure, odor and color of gills, abdominal color and vent condition parameters reached a total of 22 QI scores at the end of storage. At the beginning of storage, the total QI value was determined as 2.20±1.78. The color change in the skin of the samples according to the storage time was found to be statistically significant on the 8th day of storage and was observed as 2.00±0.70. Hemorrhage on the skin and fins was 0.80±0.44 on the 5th day of storage and 1.60±0.54 on the 8th day of storage. Eye transparency started to decrease on the 8th day of storage and was observed as completely opaque by the assessors on the 12th day of storage and completed all demerit points totaling 2. Eye pupil reached a value of 1.60±0.54 on the 8th day of storage. Eye shape started to lose its form from the 5th day of storage and reached 1.80±0.44 and 2.40±0.54 on the 5th and 8th days of storage, respectively.

Gill color and odor significantly (p<0.05) diverged from the freshness levels as of the 8th day of storage and reached 1.60 ± 0.54 and 1.80 ± 0.44 , respectively. The whole sea bass samples stored for 12 days under refrigerator conditions completed half of the total QI score on the 5th day of storage and reached a value of 11.20 ± 0.83 . Moreover, offodours were observed in the samples on the 8th day of storage and the total QI score was determined as 17.40 ± 1.94 . On the last day of storage, the samples reached a total QI score of 20.80 ± 1.30 . R²=0.97 was calculated between the QI score and storage time (Fig. 2).

FUZZY MODEL

The Fuzzy model was developed by applying a total of 12 rules to aerobically stored whole sea bass under refrigerator conditions. For the variables, 3 categories were determined as low - medium and high. According to the IF..THEN rules, the defuzzification process was applied and QI values were calculated and the categories related to sensory changes are shown in Figure 3. According to this, the categories associated with the changes of microorganisms correspond to 3 different classes in the QI score.



Figure 2. Sensory changes in the whole seabass stored under refrigerated conditions

According to the established rules, for the QI score of the "low" class, the product is consumable and meets the freshness criteria; for the score of "medium", the product begins to deviate from the first conditions in terms of freshness; the smell is pronounced, the change in color and smell of the gills is marked; and finally, for the category that matches the "high" class, the off-odors start, the eye structure has deteriorated, and the odor in the gills is prone. The corresponding variations of the calculated and observed values are shown in Figure 4. In the developed fuzzy model, the R² value is 0.96. Furthermore, the correlation between the calculated and observed QI scores is r = 0.98.

DISCUSSION

Microbiological assessment in whole sea bass stored at refrigerator conditions was investigated during 12 days of storage. Initial TMAB and TPAB counts of whole sea bass were 3.16±0.12 and 2.50±0.70 log cfu/g, respectively. Lower than the initial TMAB counts of sea bass stored under refrigerated conditions, Caklı et al.¹⁸ reported a TMAB value of 2.78±0.02 log cfu/g and a similar TPAB count of 2.47±0.06 log cfu/g in ungutted seabass stored whole on ice. In another study, Turan and Kocatepe¹⁹ investigated the effect of different modified atmosphere packaging (MAP) conditions on the shelf life of gutted seabass. The researchers reported that the initial TMAB count of gutted seabass was 2.83, and TPAB values were 3.27 log cfu/g, lower and higher than whole sea bass stored under refrigerator conditions. In a different study, MAP application effects on the microbial quality of gutted seabass were investigated. The researchers reported the initial total viable count of gutted seabass as approximately 4 log cfu/g, similar to this study.²⁰ This study's TMAB and TPAB values exceeded the microbiologi-



Figure 3. Defuzzyfication of the varibales



Figure 4. Computed vs observed values of QI scores in whole seabass stored under refrigerated temperatures

cal criterion of 6 log cfu/g²¹ on the 8th day of 12-day storage (Figure 1). The initial counts of Pseudomonas sp., LAB and members of the Enterobacteriaceae family were determined to be 3.32±0.25, 2.25±0.49 and 1.65±0.49 log cfu/g, respectively. Similar initial values for Pseudomonas sp. were reported as 3.0 log cfu/g for gutted seabass by Tsironi et al.²⁰ Higher results were reported as $4.40\pm0.12 \log cfu/g$ for whole seabass by Syropoulou et al.,²² and lower values were observed for gutted seabass by Turan and Kocatepe¹⁹ with a count of 2.52 log cfu/g. At the end of storage, Pseudomonas sp. numbers were 7.60±0.31 log cfu/g. Pseudomonas sp. is among the bacteria causing spoilage in seafood and has been reported to be dominant in the spoilage microflora of seafood.²² Parlapani et al.²³ reported Pseudomonas sp. counts in gutted seabass stored at 2°C with aerobic and MAP treatment to be 7.2 log cfu/g at the storage time when the samples were rejected at the sensory level. In this context, it is similar to the Pseudomonas sp. counts at the end

of the storage period of whole sea bass stored under refrigerator conditions. Boulares et al.24 examined the effects of LAB and citrus essential oil on the quality parameters of vacuum-packed seabass. The researchers reported that the initial LAB counts of the samples were 4.21±0.58 log cfu/g, higher than the seabass stored whole in the refrigerator, and 6.81±0.10 log cfu at the end of 21 days of storage. However, LAB counts of seabass stored whole in refrigerator conditions were determined as 4.58±0.04 log cfu/g at the end of 12 days of storage and were observed to be lower than the LAB counts reported by Boulares et al.²⁴ It has been reported that members of the Enterobacteriaceae family can grow at low temperatures but their growth is slower than that of other psychrotroph spoilers and that members of the Enterobacteriaceae family should be considered after harvesting from more polluted waters or when refrigeration or chilling is delayed.²⁵ At the end of the storage period, the number of members of the Enterobacteriaceae family was determined to be 4.27±0.14 log cfu/g in seabass stored under refrigerated conditions, which was lower than the other bacterial groups examined (Figure 1). Ucar et al.²⁶ reported the Enterobacteriaceae counts of sea bass fillets stored under cold conditions as 5.26 log cfu/g on day 6, Kritikos et al.²⁷ reported the Enterobacteriaceae counts of MAP-treated sea bass fillets stored under chilled conditions to be < 4. 6 log cfu/g, and Iacumin et al.²⁸ reported the Enterobacteriaceae counts of gutted vacuum and MAP treated sea bass kept at cooled conditions to be 4.0±0.7 and 4.5±0.8 log cfu/g on the 12th day of storage, respectively. In this context, the numbers of Enterobacteriaceae family members of sea bass stored aerobically under refrigerator conditions are similar to other studies.

QIM is one of the most used indexes to evaluate the sensory changes in seafood. The QIM developed by Knowles et al.¹⁶ for sea bass was used in this study. The QIM, which has a total of 22 demerit points (dp), was used for sea bass stored under refrigerated conditions and the changes of dp's over time are shown in <u>Figure 2</u>. The samples had an initial QI score of 2.20 ± 1.78 , indicating that the sea bass used in the study were reasonably fresh according to the QIM (0=very fresh and 17-22=spoiled, off-odour). The QI scores of the samples increased significantly (p<0.05) according to the storage time. The samples reached 11.20±0.83 on the 5th day of total storage and 20.80±1.30 at the end of storage. In the sensory evaluation of sea bass according to QIM, the assessors suggested that the samples were not consumable on the 8th day of storage (QI=17.40±1.94). In similar studies, Özogul et al.²⁹ studied sea bass fillets treated with nanoemulsion oils and stored under chilled conditions. The researchers reported that the storage time at which the samples had sensory rejection was 8 days, and the QI score was 13 in the control groups. In another study, ungutted sea bass were stored under different storage conditions (from -1°C to 22°C). The researchers reported that the samples stored in sea water (20-22 °C), storage room (16-18°C) and sea water (10-12°C) reached the sensory rejection point of 22 dp after 15, 45, and 65 hours, respectively.³⁰ In a different study, Erdağ and Ayvaz³¹ determined fish freshness in sea bass by color parameter. The researchers reported that the total QI score of 15 dp was indicated by the assessors on the 5th day of storage in sea bass kept at 4°C.

In this modelling study, microbial and sensory changes in whole sea bass stored under refrigerator conditions were investigated and the effect of microbial changes on sensory parameters during storage time was examined by applying a Fuzzy model. Moreover, by implementing a Fuzzy-based approach in which microbial parameters are used as input and sensory parameters are used as output. By developing the Fuzzy model, the shelf life of sea bass stored whole in refrigerator conditions was determined. The output values (QIM scores) calculated according to the Fuzzy model were compared with the QI scores of the samples evaluated by the assessors and the results are given in Figure 4. Accordingly, the R^2 value is 0.96 and the correlation coefficient has been determined to be *r*=0.98. There was a high correlation between the observed and calculated values. In this framework, it is evident that the developed model can predict the shelf life of whole sea bass stored in refrigerator conditions. In a study, Zare and Ghazali¹² determined the quality changes of sardine fish according to biogenic amine concentrations by applying a Fuzzy-based model. The researchers monitored the quality changes in sardines stored at different temperatures (0, 3, and 10°C) for more than 15 days. In addition, the researchers created 12 grades for the output data. According to the research results, the correlation coefficients between quality changes and storage time regarding storage temperatures were reported to be 0.97, 0.95, and 1, respectively. In another study,³² in which microbial growth parameters were used as input, the growth rates of Listeria monocytogenes were predicted according to

the Fuzzy model. The researcher created 13 Fuzzy rules using 41 growth rates. According to the rules, the correlation coefficient of the calculated and observed growth rates was 0.84. In this context, the correlation coefficients reported by the researchers are similar to the whole sea bass stored in refrigerator conditions.

CONCLUSION

In this study, microbiological spoilage of whole sea bass stored under refrigerator conditions was investigated, and microbiological and sensory indices were analyzed during shelf life in the light of sensory analysis according to QIM. In this context, according to microbiological and sensory criteria, the shelf life of sea bass fish at +4°C was determined to be 8 days. In addition, a Fuzzy model was developed in this study. In the model, a total of 12 rules were generated. The numbers of TMAB, TPAB, Pseudomonas sp., LAB, and members of the Enterobacteriaceae family were used as input. The QI scores of the samples evaluated according to the QIM method were used as output. It was observed that the Fuzzy model could determine the sensory criteria according to microbiological changes with a high correlation (r=0.98) within the 12 rules developed in this framework. Since the rules of the developed Fuzzy model apply to different seafood products, it is concluded that the model developed in this study should be investigated in other seafood products by adjusting the generated rules.

ACKNOWLEDGMENTS

The author would like to thank the assessors for attending the sensory analysis sessions.

......

AUTHOR CONTRIBUTION PER CREDIT

Conceptualization: İsmail Y. Genç (Lead). Methodology: İsmail Y. Genç (Lead). Formal Analysis: İsmail Y. Genç (Lead). Investigation: İsmail Y. Genç (Lead). Writing – original draft: İsmail Y. Genç (Lead). Writing – review & editing: İsmail Y. Genç (Lead). Funding acquisition: İsmail Y. Genç (Lead). Resources: İsmail Y. Genç (Lead). Supervision: İsmail Y. Genç (Lead).

Submitted: August 22, 2023 CST, Accepted: September 05, 2023 CST

This is an open-access article distributed under the terms of the Creative Commons Attribution 4.0 International License (CCBY-NC-ND-4.0). View this license's legal deed at https://creativecommons.org/licenses/by-nc-nd/4.0 and legal code at https://creativecommons.org/licenses/by-nc-nd/4.0/legalcode for more information.

REFERENCES

1. Grigorakis K. Compositional and organoleptic quality of farmed and wild gilthead sea bream (*Sparus aurata*) and sea bass (*Dicentrarchus labrax*) and factors affecting it: A review. *Aquaculture*. 2007;272(1-4):55-75. doi:10.1016/j.aquaculture.2007.04.062

2. Chéret R, Chapleau N, Delbarre-Ladrat C, Verrez-Bagnis V, Lamballerie MD. Effects of high pressure on texture and microstructure of sea bass (*Dicentrarchus labrax* L.) fillets. *Journal of Food Science*. 2005;70(8):e477-e483. <u>doi:10.1111/j.1365-2621.200</u> <u>5.tb11518.x</u>

3. Cakli S, Kilinc B, Cadun A, Dincer T, Tolasa S. Effects of gutting and ungutting on microbiological, chemical, and sensory properties of aquacultured sea bream (*Sparus aurata*) and sea bass (*Dicentrarchus labrax*) stored in ice. *Critical Reviews in Food Science and Nutrition*. 2006;46(7):519-527. doi:10.1080/10408 390500343209

4. Pinto de Rezende L, Barbosa J, Teixeira P. Analysis of alternative shelf life-extending protocols and their effect on the preservation of seafood products. *Foods*. 2022;11(8):1100. doi:10.3390/foods11081100

5. Gram L. Microbiological spoilage of fish and seafood products. In: *Compendium of the Microbiological Spoilage of Foods and Beverages*. Springer; 2009:87-119. <u>doi:10.1007/978-1-4419-082</u> <u>6-1_4</u>

6. Levin RE. Assessment of seafood spoilage and the microorganisms involved. In: *Handbook of Seafood and Seafood Products Analysis*. CRC Press; 2009:515-535. doi:10.1201/9781420046359-c28

7. Odeyemi OA, Alegbeleye OO, Strateva M, Stratev D. Understanding spoilage microbial community and spoilage mechanisms in foods of animal origin. *Comp Rev Food Sci Food Safe*. 2020;19(2):311-331. <u>doi:10.11</u> <u>11/1541-4337.12526</u>

8. Martinsdóttir E, Schelvis R, Hyldig G, Sveinsdóttir K. Sensory evaluation of seafood: methods. In: *Fishery Products: Quality, Safety and Authenticity*. Wiley-Blackwell; 2009:425-443. <u>doi:10.1002/9781444</u> 322668.ch20

9. Esteves E, Aníbal J. Sensory evaluation of seafood freshness using the quality index method: A metaanalysis. *International Journal of Food Microbiology*. 2021;337:108934. <u>doi:10.1016/j.ijfoodmicro.2020.108</u> <u>934</u> 10. Mejlholm O, Dalgaard P. Modeling and predicting the growth boundary of *Listeria monocytogenes* in lightly preserved seafood. *Journal of Food Protection*. 2007;70(1):70-84. <u>doi:10.4315/0362-028x-70.1.70</u>

11. Lalabadi HM, Sadeghi M, Mireei SA. Fish freshness categorization from eyes and gills color features using multi-class artificial neural network and support vector machines. *Aquacultural Engineering*. 2020;90:102076. doi:10.1016/j.aquaeng.2 020.102076

12. Zare D, Ghazali HM. Assessing the quality of sardine based on biogenic amines using a fuzzy logic model. *Food Chemistry*. 2017;221:936-943. <u>doi:10.101</u> 6/j.foodchem.2016.11.071

13. Moosavi-Nasab M, Khoshnoudi-Nia S. Combining knowledge-and data-driven fuzzy approach to evaluate shelf-life of various seafood products. *Food Quality and Safety*. 2021;5:fyab022. <u>doi:10.1093/fqsaf</u> <u>e/fyab022</u>

14. Vivek K, Subbarao KV, Routray W, Kamini NR, Dash KK. Application of fuzzy logic in sensory evaluation of food products: A comprehensive study. *Food Bioprocess Technol*. 2020;13(1):1-29. <u>doi:10.100</u> 7/s11947-019-02337-4

15. Syropoulou F, Parlapani FF, Anagnostopoulos DA, Stamatiou A, Mallouchos A, Boziaris IS. Spoilage Investigation of Chill Stored Meagre (*Argyrosomus regius*) Using Modern Microbiological and Analytical Techniques. *Foods*. 2021;10(12):3109. <u>doi:10.3390/foo</u> <u>ds10123109</u>

16. Knowles TG, Brown SN, Warriss PD, et al. Effect of electrical stunning at slaughter on the carcass, flesh and eating quality of farmed sea bass (*Dicentrarchus labrax*). *Aquaculture Res.* 2007;38(16):1732-1741. do i:10.1111/j.1365-2109.2007.01846.x

17. Zadeh LA. A computational approach to fuzzy quantifiers in natural languages. In: *Computational Linguistics*. Pergamon; 1983:149-184. <u>doi:10.1016/b97</u> <u>8-0-08-030253-9.50016-0</u>

18. Cakli S, Kilinc B, Cadun A, Dincer T, Tolasa S. Quality differences of whole ungutted sea bream (*Sparus aurata*) and sea bass (*Dicentrarchus labrax*) while stored in ice. *Food Control*. 2007;18(5):391-397. doi:10.1016/j.foodcont.2005.11.005

19. Turan H, Kocatepe D. Different MAP conditions to improve the shelf life of sea bass. *Food Sci Biotechnol*. 2013;22(6):1589-1599. <u>doi:10.1007/s10068-013-025</u> <u>5-x</u>

20. Tsironi T, Ntzimani A, Gogou E, et al. Modeling the effect of active modified atmosphere packaging on the microbial stability and shelf life of gutted sea bass. *Applied Sciences*. 2019;9(23):5019. <u>doi:10.3390/a pp9235019</u>

21. ICMSF. International Commission on Microbiological Specifications for Foods. Sampling plans for fish and shellfish. In: ICMSF, ed. *Microorganisms in Food. Sampling for Microbiological Analysis: Principles and Scientific Applications*. University of Toronto Press; 1992.

22. Syropoulou F, Parlapani FF, Kakasis S, Nychas GJE, Boziaris IS. Primary processing and storage affect the dominant microbiota of fresh and chill-stored sea bass products. *Foods*. 2021;10(3):671. <u>doi:1</u> 0.3390/foods10030671

23. Parlapani FF, Haroutounian SA, Nychas GJE, Boziaris IS. Microbiological spoilage and volatiles production of gutted European sea bass stored under air and commercial modified atmosphere package at 2°C. *Food Microbiology*. 2015;50:44-53. <u>doi:10.1016/j.f</u> <u>m.2015.03.006</u>

24. Boulares M, Ben Moussa O, Mankai M, Sadok S, Hassouna M. Effects of lactic acid bacteria and citrus essential oil on the quality of vacuum-packed sea bass (*Dicentrarchus labrax*) fillets during refrigerated storage. *Journal of Aquatic Food Product Technology*. 2018;27(6):698-711. doi:10.1080/10498850.2018.1484 544

25. Papadopoulos V, Chouliara I, Badeka A, Savvaidis IN, Kontominas MG. Effect of gutting on microbiological, chemical, and sensory properties of aquacultured sea bass (*Dicentrarchus labrax*) stored in ice. *Food Microbiology*. 2003;20(4):411-420. <u>doi:10.10</u> <u>16/s0740-0020(02)00148-x</u>

26. Ucar Y, Özogul Y, Özogul F, Durmuş M, Köşker AR. Effect of nisin on the shelf life of sea bass (*Dicentrarchus labrax* L.) fillets stored at chilled temperature (4±2°C). *Aquacult Int.* 2020;28(2):851-863. doi:10.1007/s10499-020-00512-5 27. Kritikos A, Aska I, Ekonomou S, et al. Volatilome of chill-stored european seabass (*Dicentrarchus labrax*) fillets and atlantic salmon (*Salmo salar*) slices under modified atmosphere packaging. *Molecules*. 2020;25(8):1981. doi:10.3390/molecules25081981

28. Iacumin L, Jayasinghe AS, Pellegrini M, Comi G. Evaluation of Different Techniques, including Modified Atmosphere, under Vacuum Packaging, Washing, and *Latilactobacillus sakei* as a Bioprotective Agent, to Increase the Shelf-Life of Fresh Gutted Sea Bass (*Dicentrarchus labrax*) and Sea Bream (*Sparus aurata*) Stored at 6 ± 2 °C. *Biology*. 2022;11(2):217. do i:10.3390/biology11020217

29. Özogul Y, Durmus M, Ucar Y, Özogul F, Regenstein JM. Comparative study of nanoemulsions based on commercial oils (sunflower, canola, corn, olive, soybean, and hazelnut oils): Effect on microbial, sensory, and chemical qualities of refrigerated farmed sea bass. *Innovative Food Science* & Emerging Technologies. 2016;33:422-430. doi:10.10 16/j.ifset.2015.12.018

30. Šimat V, Soldo A, Maršić-Lučić J, Tudor M, Bogdanović T. Effect of different storage conditions on the dielectric properties of the sea bass (*Dicentrarchus labrax*, L.). *Acta Adriatica: International Journal of Marine Sciences*. 2009;50(1):5-10. <u>https://hr</u> <u>cak.srce.hr/40264</u>

31. Erdağ M, Ayvaz Z. The Use of Color to Determine Fish Freshness: European Seabass (*Dicentrarchus labrax*). Journal of Aquatic Food Product Technology. 2021;30(7):847-867. <u>doi:10.1080/10498850.2021.1949</u> 771

32. Genç İY. Prediction The Growth Rate Of *Listeria monocytogenes* Based On Fuzzy Logic. In: 2nd *International Congress of the Turkish Journal of Aqriculture – Food Science and Technology*. KKTC; 2021:978-605-80461-4-6.