

## Original Research Articles

# The presence of microplastics affects *Spiella japonica* hatching performance and microbiota colonization

Tao Zhang<sup>1,2a</sup>, Xinya Chen<sup>3b</sup>, Ke Sun<sup>3</sup>, Jiyi Chen<sup>3</sup>, Xinxin Chen<sup>3</sup>, Jilin Xu<sup>1</sup>, Huilai Shi<sup>2</sup>, Hongling Ping<sup>2</sup>, Jie He<sup>2</sup>, Bin Li<sup>2</sup>, Yebing Yu<sup>3</sup>, Lei Li<sup>4</sup>, Yanming Sui<sup>3c</sup>

<sup>1</sup> School of Marine Sciences, Ningbo University, Ningbo, China, <sup>2</sup> Zhejiang Province Key Lab of Mariculture and Enhancement, Zhejiang Marine Fisheries Research Institute, Zhoushan, China, <sup>3</sup> College of Marine and Biology Engineering, Yancheng Institute of Technology, Yancheng, China, <sup>4</sup> East China Sea Fisheries Research Institute, Chinese Academy of Fishery Sciences, Shanghai, China

Keywords: Microplastics, *Sepiella maindroni*, hatching performance, 16S rRNA, microflora

<https://doi.org/10.46989/001c.94204>

## Israeli Journal of Aquaculture - Bamidgah

Vol. 76, Issue 1, 2024

Microplastics (MPs) are a global concern regarding environmental pollution. This study evaluated the impacts of MPs with two sizes (5 µm and 0.5 µm) on hatching performance and microbiota of *Sepiella japonica*. The presence of MPs increased the hatching rate at some stages of the fertilization process and reduced the oxygen consumption rate at the gastrula stage. No size-dependent impact was observed. The 16S rRNA gene was sequenced to identify the flora. Clustering tags assessed species diversity in the samples with 97% similarity. Proteobacteria was the most abundant phylum in all three groups. MPs publicity appreciably modified flower structure. The most variable genera were *Ralstonia*, *Methylophilus*, and *Pseudorhodoferrax*, which can furnish nutrients and modify the host's immune response. MPs exposure appeared to enrich beneficial bacteria in this study. The presence of MPs with a size of 5µm played a greater role in this process, which is supported by presumptive functions. However, since the adsorption of suspended MPs on aquatic eggs can have cascading effects on specific life stages of oviparous animals, regular monitoring of microbial communities is necessary after juvenile *S. japonica* formation to prevent disease outbreaks.

## INTRODUCTION

Many invertebrates and vertebrates, including chordates, arthropods, echinoderms, cnidarians, and others, lay eggs that develop ex vivo in the water environment. These envelopes are simple coats in cnidarians and nematodes, but they are acellular membranes with or without an external jelly coating in chordates, arthropods, echinoderms, and mollusks.<sup>1</sup> In addition to the inner vitelline membrane, teleost fish eggs, pterygoid insect eggs, and cephalopod eggs are all covered by an external chorion membrane. Chorion and vitelline membranes cooperate to protect growing embryos from external pressures in the water column, including mechanical forces, predators, and infections.<sup>2</sup>

According to studies on teleost fish eggs, the chorion functions as a reliable physical barrier against the ingestion of particle pollutants. For instance, it was discovered that microplastics (MPs) particles adhered to the villi on the surface of chorion in medaka fish eggs but were absent in

the eggs.<sup>3</sup> Since cuttlefish are cephalopods, they lack villi, and a recent analysis discovered MPs in the cuttlefish embryo.<sup>4</sup>

In the last few decades, the production of plastics has increased significantly, from 500 to 350,000 tons annually.<sup>5</sup> In all seas, including the pelagic, abyssal, and polar zones, over 10% of this plastic pollution makes its way into marine environments, and the accumulation of plastic debris becomes a severe issue.<sup>6</sup> More than 92% of the plastic waste in the ocean nowadays is made up of MPs, described as polymer particles typically having a diameter of less than 5 mm. They are mostly made from manufactured plastic beads or the fragmentation of larger plastics.<sup>7</sup> Recent estimates place the amount of MPs particles in the world's oceans at between 15 and 51 trillion, or 93,000 and 236,000 tons.<sup>8</sup> As reported, there may be more than 100,000 microplastic particles per cubic meter in coastal waters.<sup>9</sup>

Cuttlefish *Sepiella japonica* is an endemic local species in the East China Sea.<sup>10</sup> Breeding reproduction is essential for this species because of its excessive consumption and

a Author equally contributed to the study

b Author equally contributed to the study

c Corresponding Author: Yanming Sui, Email: 2181642148@qq.com

ecological deterioration.<sup>11,12</sup> MPs suspended in the surface water of the Yangtze estuary, which is the major habitat for the mariculture of *S. japonica*, can reach up to  $4137.3 \pm 2461.5$  n/m<sup>3</sup>.<sup>13</sup> Fertilized eggs are deposited on or adhered to the coral reef from April to June. Seawater is used to soak the fertilized eggs. The juveniles promptly emerge and have typical swimming abilities after a month of in vitro incubation. It was the most vulnerable during the hatching period. The marine environment's dissolved oxygen, pH, temperature, and light had little effect.<sup>14</sup> Although there is no solid evidence that MPs can stick to *S. japonica* eggs, a study on zebrafish eggs with a similar structure revealed that MPs of various sizes in the seawater performed differently when connected to the chorion surface. Smaller MPs of 1 to 5  $\mu$ m can cover the entire chorion surface, but relatively large MPs of 10 to 20  $\mu$ m are only attached in patches.<sup>15</sup> MPs have been proven to obstruct the egg's pores, which is likely to minimize the supply of oxygen to the embryo, resulting in hypoxia and the embryo hatching postponed.<sup>16</sup> In order to benefit both the host and themselves, these symbiotic bacteria release some compounds that the host needs for growth and development in addition to using the nutrients from the host. Both the host and parasitic bacteria adapt to changes in the seawater environment. According to studies, MPs exposure can alter the intestinal microbiota of marine Organisms, increasing the abundance of Firmicutes and decreasing the abundance of Bacteroides.<sup>17-19</sup> However, the response of the bacterial community surrounding the fertilized eggs of *S. japonica* to the presence of marine MPs is still unavailable. Thus, in this study, we quantify the emergence rate, emergence cycle, and deformity rate of the fertilized eggs of *S. japonica*; physiological indicators such as respiratory oxygen consumption rate and ammonia nitrogen excretion rate of the fertilized eggs of *S. japonica*; to reveal the impact of MPs exposure on the diversity of bacteria on the fertilized eggs of *S. japonica* at the genumacro genome using macro genome sequencing to explore the potential impact of MPs on *S. japonica*. The findings will yield evidence that can be used to shape future regulations surrounding the hazards of marine MPs pollution.

## 2. MATERIALS AND METHODS

### 2.1. CUTTLEFISH PREPARATION

The experiment was conducted from October 2020 to August 2021. The *S. japonica* eggs were collected from the parent species' natural broods in an indoor cement pond that was maintained by an aquatic breeding facility in Zhoushan, Zhejiang Province. The parents were lowered into the cement pool (6 m×4 m×1.4 m) with carcasses measuring (11.5–12.4 cm in length, 7.5–8.2 cm in breadth, and 109.5–159.4 g in body weight). Natural seawater with a salinity of 24.4–28.2, a water temperature of 19–21 °C, a pH of 7.38–8.21, a COD of 5.45–7.5 mg/L, continuous aeration, and a 50% water daily change is used for parent breeding and egg incubation.

### 2.2. SEAWATER PREPARATION

Nine tanks of 50 L in total (triplicates for each of the three MPs treatments: Control, 5  $\mu$ m, and 0.5  $\mu$ m) were used in our investigation. In this investigation, commercial microspheres (ACMETM, 5  $\mu$ m polystyrene (PS),  $3.6 \times 10^8$  items/mL, and 0.5  $\mu$ m PS,  $3.6 \times 10^{11}$  items/mL) were used to prepare the MPs suspensions. According to previous research, the MPs concentration was set to  $10^4$  items/L, representing the concentration at the primary location for *S. japonica* culture.<sup>20</sup> Airflow was performed throughout the experiment to keep the microsphere particle suspensions in place.

### 2.3. EFFECTS OF MPS ON THE EARLY DEVELOPMENT OF *S. JAPONICA*

For spawning, the air-exposure techniques linked to flow stimulation were used. In a nutshell, 20 participants were moved to the 0.3 m<sup>2</sup> square tank after 4 hours of air exposure. A submersible pump maintained an emergent flow of seawater. Eggs were collected using a 400-mesh net for 10 minutes. The eggs were transferred to the tanks containing the right seawater for the cuttlefish hatching and development after being rinsed in the appropriate seawater (15 eggs/mL). Each tank's 3×100 eggs were sampled at four different developmental stages: the gastrula stage (GS), the organization formed stage (OF), the endoskeleton formation stage (EFS), and the embryo just before hatching (EBH). The samples were then stored with 4% paraformaldehyde for further study. The standard for morphological classification was displayed in [Figure 1](#). Most fertilized eggs developed to the morphological criteria in [Figure 1](#) for stage time classification.


The hatching rate (%) was calculated by samples taken at 4 stages after fertilization.

Hatching rate = (the fertilized eggs meet the morphological classification standard) /  $100 \times 100^{21}$  is referred to morphological classification when determining whether larvae have a distorted or normal shape.

### 2.4. EFFECTS OF MPS ON PHYSIOLOGICAL INDEXES OF *S. JAPONICA*

For the test to measure oxygen intake and ammonia excretion rate at 4 stages, a conical flask with a volume of 500 mL was filled with saltwater after sand filtering. The bottle-neck was sealed with paper film after 100 fertilized eggs or embryos had been placed within. Three parallels were set for each treatment, and the test was run at  $(20 \pm 1)$  °C. The eggless conical flask was used as a placeholder. The examination lasted for six hours.

Oxygen consumption rate measurement: Dissolved oxygen (DO) was measured using WINKLER iodometry. The oxygen consumption rate is calculated as  $OR = [(DO_t - DO_0) V] / (Nt)$ . Where:  $DO_t - DO_0$  in the respiratory bottle at the test's end, mg/L;  $DO_0 - DO_0$  in the blank bottle at the test's finish, mg/L; OR- oxygen consumption rate per egg, mg/(ind.h); V stands for the volume of water in the conical flask, N for the number of fertilized eggs in the bottle, T for the length of the test, h.



**Figure 1. The morphological classification standard for the fertilized egg development stage of *Sepiella japonica*, A: gastrula stage (GS), B: organization formed (OF), C : endoskeleton formation stage (EFS), D: the embryo before hatching (EBH).**

The ammonia excretion (AN) was calculated using the SKALARSAN++interval flow analyzer. Ammonia rate calculation:  $NR = [(AN_0 - AN_t) V] / (Nt)$ . Where:  $AN_t$ -AN in the respiratory bottle at the conclusion of the test, mg/L;  $AN_0$ -AN in the blank bottle at the end of the test, mg/L; NR-ammonia excretion rate per egg, mg/(ind.h); V stands for the volume of water in the conical flask, N for the number of fertilized eggs in the bottle, T for the length of the test, h.


## 2.5. 16S RNA GENE SEQUENCING AND ANALYSIS

35-40 eggs were randomly selected from each tank at the embryonic stage to create 9 samples for 16S RNA gene sequencing and further analyses. Total genomic DNA was extracted using a DNA Extraction Kit and following the manufacturer's instructions. NanoDrop and an agarose gel were used to confirm the DNA's purity and amount. To be processed further, the extracted DNA was diluted to a concentration of 1 ng/L and kept at -20°C. Using the barcoded primers and Takara Ex Taq, the diluted DNA was employed as a template for the polymerase chain reaction (PCR) amplification of the bacterial 16S ribosomal ribonucleic acid (rRNA) genes (Takara). V3-V4 (or V4-V5) variable portions of 16S rRNA genes were amplified for bacterial diversity investigation using universal primers 343F and 798R (or 515F and 907R for V4-V5 regions). Variable portions of the 18S rRNA gene were amplified using the universal primers 817F

and 1196R to investigate Eukaryota diversity. ITS I variable areas were amplified to investigate fungal diversity using the universal primers ITS1F and ITS2.

Using gel electrophoresis to assess amplicon quality, AMPure XP beads (Agencourt) were used to purify the amplicons before being amplified for a subsequent cycle of PCR. The final amplicon was measured using a Qubit ds-DNA assay kit after purification using AMPure XP beads. For further sequencing, identical amounts of purified amplicons were pooled.

FASTQ format was used for the raw sequencing data. Using Trimmomatic software,<sup>22</sup> paired-end readings were preprocessed to identify and remove ambiguous bases (N). Low-quality sequences with an average quality score of 20 were also terminated using a sliding window trimming method. Using FLASH software, paired-end readings were assembled after cutting.<sup>23</sup> Ten base pairs (bp) of least overlap, 200 bp of maximum overlap, and a maximum misfit rate of 20% made up the assembly parameters. To further denoise the sequences, reads with base pairs shorter than 200 bp or ambiguous or homologous sequences were eliminated, whereas reads with 75% of bases higher than Q20 were kept. Chimera-containing reads were then found and eliminated. Using the QIIME software, these two actions were completed (version 1.8.0).<sup>24</sup> Using V search software with a 97% similarity criterion, clean reads were subjected to primer sequence removal and clustering to produce op-



**Figure 2.** Hatching rate (HR) of *S. japonica* exposed to microplastics with different sizes: Control, 0.5µm and 5µm during six days. Each day, the symbols denoted by different lowercase letters mean HR significantly differs among the three microplastic treatments ( $P < 0.05$ ).

erational taxonomic units (OTUs).<sup>25</sup> With the help of the QIIME program, the representative read for each OTU was chosen. The Silva database Version 123 (or Greengenes) (16S rDNA) was blasted against using an RDP classifier, which had a 70% confidence threshold, and all representative reads were annotated.<sup>26</sup> Using Blast, the Unite database (ITSs rDNA) was also searched for all representative reads.<sup>27</sup>


## 2.6. STATISTICAL ANALYSES

This study presented the results as Mean  $\pm$  Standard Deviation (SD). Shapiro-test Wilk's and Levene's test (both using SPSS 26.0) were used to determine the normality and homogeneity of the data. One-way ANOVA was used to examine the significant effects of MPs for each treatment, and Tukey's HSD test was used for multiple comparisons. Molecular variance (AMOVA) analysis was used to evaluate bacterial communities. The threshold for a significant difference was defined as a  $P$  value of less than 0.05.

## 3. RESULTS AND DISCUSSION

The results indicated that  $10^4$  items/L PS (0.5 µm and 5 µm) may improve the hatching rate when compared to the control during the hatching process ( $P < 0.05$ ), but there was no significant difference at the end of the hatching process ( $P > 0.05$ , [Figure 2](#)). There was no significant difference in the deformation rate across the treatments ( $P > 0.05$ , [Figure 3](#)).


These findings differed from those of several previous reports,<sup>28-30</sup> which suggested that MPs can decrease hatch-



**Figure 3.** Deformity rate (DR) of *S. japonica* exposed to microplastics with different sizes

ing success, delay the hatching process, and increase the deformity rate. It is believed that the adsorption of fine particles onto eggshells in the presence of dense MPs could obstruct spiracles and channels, resulting in a decrease in the absorption of oxygen and nutrients. Fertilized eggs may experience a longer hatching period.<sup>16</sup> These studies primarily focused on teleost fish. Additionally, several investigations revealed that MPs had little impact on the development of fertilized eggs.<sup>31,32</sup> Additionally,<sup>33</sup> discovered that low levels of polystyrene MPs reduce marine medaka's early developmental toxicity to phenanthrene (*Oryzias melastigma*). To the best of our knowledge, the metabolic





**Figure 4.** Oxygen consumption rate (OR) of *S. japonica* exposed to microplastics with different sizes at four developmental stages. The column denoted by different lowercase letters at each developmental stage means OR significantly differs among the three microplastic treatments ( $P < 0.05$ ). The column denoted by different capital letters under each microplastic treatment means OR significantly differs among the four developmental stages ( $P < 0.05$ ).

reactions to MPs from marine bivalves varied depending on the MPs' characteristics (including kind, shape, and size), as well as the exposure modes (including length and abundance).<sup>34</sup> In contrast,<sup>15</sup> size-dependent effects of MPs were not confirmed in this study. Species-specific differences in hatching performance could exist. Therefore, further research on various species needs to be done.

At the GS stage, MPs were observed to have a significantly decreased oxygen consumption rate ( $P < 0.05$ , [Figure 4](#)), suggesting that MPs may have blocked the spiracles of *S. japonica* fertilized eggs at the start of development. However, the significant difference vanished from the OF stage, demonstrating that the negative effects of MPs were transient. Under three treatments, the oxygen consumption rate often rose as development lengthened. Furthermore, because organisms would increase their ammonia excretion rates to combat environmental stress,<sup>35</sup> there was no significant difference in ammonia excretion rates between treatments ( $P > 0.05$ , [Figure 5](#)), implying that  $10^4$  items/L PS ( $0.5 \mu\text{m}$  and  $5 \mu\text{m}$ ) had no negative effects on *S. japonica* fertilized eggs.

On an Illumina MiSeq platform, 9 samples produced excellent paired-end reads (200 bp). 676,841 contigs were created by preprocessing and assembling the raw data. The average length of the effective tags ranged from 63,279 to 68,241, was 387–437 nt, making up 85.72%–90.74% of the raw data. The effective tags were clustered into OTUs at 97% similarity to evaluate the species diversity within samples. 9 samples yielded 1553 OTUs, ranging in size from 223 to 446.

Distribution histograms created using a maximum sorting method display the top 15 most prevalent microorganisms at various taxonomic levels (phylum, class, order, family, genus, and species). The maximum number of organisms in each phylum in all samples was determined, and the top 15 phyla were chosen for the figure. Proteobacteria, Bacteroidota, Firmicutes, Actinobacteria, Desulfobacterota, Gemmatimonadota, Verrucomicrotiota, Bdellovibrionota, Acidobacteriota, Patescibacteria, Deferribacterota, Campilobacterota, Fusobacteriota, and Chloroflexi were the top 15 ([Figure 6](#)). *Tenacibaculum*, *Muribaculaceae*, *Ruegeria*, *Nautella*, *Roseovarius*, *Leisingera*, *Ralstonia*, *Acinetobacter*, *Vibrio*, *Bacteroides*, *Methylophilus*, *Azospirillum*, *Escherichia-Shigella*, *Kordiimonas*, and *Methylobacillus* were the most prevalent genera ([Figure 7](#)).

By calculating the abundance of each OTU ([Figure 8](#)), choosing the top 50 OTUs based on their abundance of tags, and creating an evolution tree, it was possible to discover the phylogenetic relationships between various OTUs and the variations among the dominant species. Proteobacteria, Bacteroidota, Actinobacteriota, Firmicutes, Deferribacterota, Myxococcota, and Verrucomicrobiota were the taxa with the highest OUT diversity.

The dominating microbiotas in the three groups were examined by normalizing the OTU table and examining both the shared and unique OTUs on the microbial community compositions of cuttlefish fertilized eggs at varied PS sizes. The OTUs in the digestive tracts were identified using a flower plot diagram ([Figure 9](#)). A total of 255 OTUs were clustered into the 4 groups. The OTUs 850, 761, and 708




Figure 5. Ammonia excretion rate (AE) of *S. japonica* exposed to microplastics with different sizes at four developmental stages: gastrula stage (GS), organization formed (OF), endoskeleton formation stage (EFS), the embryo before hatching (EBH).




Figure 6. Distribution of the top 15 phyla in each sample. cd (1–3), c5 $\mu\text{m}$  (1–3) and c500nm (1–3) represent the contents at different treatment conditions.




Figure 7. Distribution of the top 15 phyla in each sample. cd (1–3), c5 $\mu$ m (1–3) and c500nm (1–3) represent the contents at different treatment conditions.





Figure 8. Phylogenetic relationship of OTUs. On the left is a phylogenetic tree graph, and phylum is the gate information; on the right is an abundance graph corresponding to the abundance of the left OTU in each sample.




**Figure 9.** OTUs distribution flower plot. The numbers in the Core in the picture represent the common OTUs in all the samples (Core OTUs), and the numbers on the petals represent the total OTUs of each sample minus the number of common OTUs.

were specific to the c500nm (1-3) wavelength, while the OTUs 1181, 817, and 749 were specific to the control (cd) (1-3), and the OTUs 751, 758, and 812 were specific to c5µm (1-3).

The top 15 genus-level microorganisms were most prevalent in each treatment group (Figure 10). *Ruegeria*, *Nautella*, *Roseovarius*, and *Leisingera* were the top four prevalent genera among the samples in the cd groups. *Muribaculaceae*, *Acinetobacter*, *Vibrio*, *Bacteroides*, *Methylophilus*, *Azospirillum*, *Escherichia-Shigella*, and *Methylobacillus* were the most prevalent genera in the samples from cd groups c5m. *Ralstonia* was the most prevalent genus in the cd groups c500nm samples. Based on the relative abundances of OTUs, the weighted UniFrac distances of all nine samples within the four treatment groups were determined. Utilizing the weighted UniFrac distance, we conducted principal coordinate analysis (PCoA) (Figure 11). All of the samples, including all groups, did not cluster properly and were not arranged along the chart's first main component (PC1). They accounted for 73.58% of the total weighted PCoA. A sample clustering tree was created using the unweighted pair group technique with arithmetic mean (UPGMA) to examine further the similarities between various samples and groups (Figure 12). It was clear that there

was no visible clustering of the groups or samples into a single branch, which was in line with the PCoA analysis. The variations in bacterial communities across the four groups were then further evaluated using an analysis of molecular variance (AMOVA). The cd and c5µm groups of communities in *Ralstonia*, *Vibrio*, *Bacteroides*, *Methylophilus*, *Methylobacillus*, *Methylovorus*, *Demequina*, *Pseudorhodofera*, and *Comamonas* showed distinct characteristics. There were clear variances between the cd and c500nm groups of communities in *Ralstonia*, *Methylophilus*, *Methylovorus*, *Demequina*, *Pseudorhodofera*, *Comamonas*, and *Bradyrhizobium*. The cd and c5µm groups of communities in *Ralstonia*, *Vibrio*, *Bacteroides*, *Methylophilus*, *Methylobacillus*, *Methylovorus*, *Demequina*, *Pseudorhodofera*, and *Comamonas* showed distinct characteristics. There were clear variances between the cd and c500nm groups of communities in *Ralstonia*, *Methylophilus*, *Methylovorus*, *Demequina*, *Pseudorhodofera*, *Comamonas*, and *Bradyrhizobium*. Significant differences existed between the c5 µm and c500nm groups of communities in *Vibrio*, *Bacteroides*, *Methylophilus*, *Pseudorhodofera*, and *Bradyrhizobium*. (Figure 13). All three findings demonstrated that the microbial community structure of





**Figure 10. The top 15 genera based on the relative abundance in the group's cd, c5µm, and c500nm at the genus classification level represent the contents at different treatment conditions.**

the cuttlefish fertilized eggs is significantly impacted by variations in MPs size.

We performed LEfSe on taxa while keeping as many as possible for useful comparisons in order to further discover significant differences in biomarkers among the categories. Only six taxa, comprising the genera *Agaribacterium*, *Halioxenophilus*, *Aquimarina*, *Dechloromonas*, *Limibaculum*, and *Flavonifractor*, were found to be more prevalent in group cd, according to the comparison. Group c5µm, however, has a higher proportion of eleven taxa, including five NB genera, *Methylophilaceae*, *Comamonadaceae*, *Sandaracinaceae*, *Methylophilus*, *Pseudorhodoferrax*, and *Koukoulia* (Figure 14). When seven taxa, including the genera *Burkholderiaceae*, *Ceadibacteraceae*, *Ralstonia*, *Bradyrhizobium*, *Comamonas*, *Phreatobacter*, and *Caedibacterales*, were more prevalent in the c500nm group.

By utilizing PICRUST to anticipate metagenomes, the presumed roles of the cuttlefish fertilized egg microbiome were examined. Figure 15 displays the top seven enrichment pathways: Genetic information processing is listed first, followed by unclassified, cellular processes, organismal systems, human diseases, metabolism, and finally, unclassified. There was no obvious major enrichment pathway. In relation to the seven paths, Group c5µm performs mediocly. Meanwhile, Group cd functions heavily in the road for human diseases, and Group c500nm functions strongly in the pathway for cellular processes and organismal systems.

The findings demonstrated that Proteobacteria dominated the phylum flora in cuttlefish fertilized eggs and that MPs tended to lower Proteobacteria's relative abundance.


According to,<sup>36-38</sup> proteobacteria were also found to be prevalent in the Pacific oyster *Crassostrea gigas*, noble scallop *Chlamys nobilis*, and Yesso scallop *Patinopecten yessoensis*, indicating their biological and ecological significance in the mollusk. Proteobacteria would become less numerous under environmental stress, according to the discovery that the relative abundance of these microorganisms declined steadily under high-temperature stress.<sup>37</sup> The outcomes are consistent with this investigation. Healthy individuals' intestines also include symbiotic Proteobacteria; nevertheless, they may develop into intestinal microorganisms that cause inflammation in certain intestinal environments. Intestinal Proteobacteria can represent microecological problems or unstable intestinal microbial community structures.<sup>39</sup> Proteobacteria in the digestive tract of marine mollusks have received the majority of attention in pertinent studies to date. Its role and existence in other organs require more research.

Three genera, *Ralstonia*, *Methylophilus*, and *Pseudorhodoferrax*, have significantly varied relative abundances among the three treatment groups. We hypothesize that MPs alter the composition of the microflora community in the fertilized egg of *S. japonica*. Zebrafish eggs, which have an exterior structure resembling *S. japonica*, can adsorb micro- and nanoparticles.<sup>40</sup> Adsorbed particles may promote microbial diversity and have antibacterial effects, which closely agrees with our results. A species from *Ralstonia* with high amounts of poly-hydroxybutyrate, according to,<sup>41</sup> can control the immunological response in mussel larvae exposed to *Vibrio coralliilyticus*. *Methylophilus* is believed to contribute significant nutrients to an algal diet.<sup>42</sup> Catalase can be produced by pseudorhodoferrax.<sup>43</sup> This study's MPs exposure appears to enhance the good bacteria, as shown by an increase in the hatch rate during fertilization. However, the adsorption of dispersed antimicrobial nanoparticles on aquatic eggs may have cascade effects on several oviparous animal life stages, according to.<sup>40</sup> According to one study, MPs could interact with microorganisms directly and indirectly by acting as a carrier for other contaminants.<sup>44</sup> After the development of the *S. japonica* juveniles, ongoing surveillance of microbial organisms is required to stop the spread of the disease.

The microbial makeup of the mussel digestive system was investigated in the current work using 16S rRNA second-generation sequencing in response to variations in exposure to MPs of various sizes. We discovered that exposure to MPs impacted a number of dominating bacteria. The findings of the current study may serve to illuminate further the possible effects of growing marine MPs exposure on cuttlefish fertilized eggs. This information should be crucial for addressing the threats that marine MPs pollution poses to the cuttlefish breeding sector.

#### CONFLICT OF INTEREST

*The authors declare that the research was conducted without any commercial or financial relationships that could be construed as a potential conflict of interest.*



**Figure 11. PCoA scores are based on the relative abundances of OTUs in the 9 samples (based on the weighted UniFrac distance).**


#### FUNDING

This study was funded by research grants from the National Natural Science Foundation of China (No.41706142 & 32071615), the Open Project Fund of Zhejiang Province Key Lab of Mariculture and Enhancement (2021KF003), and Funding for school-level research projects of Yancheng Institute of Technology (XJR2021035). Yanming Sui is supported by a fellowship from China Scholarship Council.


#### AUTHORS' CONTRIBUTION PER CREDIT

Conceptualization: Tao Zhang (Lead). Project administration: Xinya Chen (Lead). Supervision: Ke Sun (Lead). Formal Analysis: Jiyi Chen (Lead). Data curation: Xinxin Chen (Lead). Software: Jilin Xu (Lead). Validation: Huilai Shi (Lead). Visualization: Hongling Ping (Lead). Resources: Jie He (Lead). Methodology: Bin Li (Lead). Writing – original draft: Yebing Yu (Lead). Investigation: Lei Li (Lead). Writing – review & editing: Yanming Sui (Lead).


Submitted: November 17, 2023 CST. Accepted: December 31, 2023 CST. Published: February 28, 2024 CST.



**Figure 12.** UPGMA cluster phylum level tree based on the weighted UniFrac distance. cd (1–3), c5um (1–3), and c500nm (1–3) represent the contents at different treatment conditions.



**Figure 13.** Perform statistical analysis at the genus level, select the relative abundance of Top 10 and do a relative abundance boxplot analysis to quickly obtain the abundance of the dominant differential species within the group and compare between groups.



**Figure 14.** Bacterial taxa differentially represented among the three groups as determined by LEfSe using default parameters




Figure 15. Predicted functions of microbial floras in the esophagus and intestine. cd (1–3), c5µm (1–3), and c500nm (1–3) represent the contents at different treatment conditions. Calculate the predicted KEGG results at three levels based on the Kruskal–Wallis algorithm to calculate the difference. If there is no difference or the number of differences is less than 2, cannot make a map.



This is an open-access article distributed under the terms of the Creative Commons Attribution 4.0 International License (CCBY-4.0). View this license's legal deed at <http://creativecommons.org/licenses/by/4.0> and legal code at <http://creativecommons.org/licenses/by/4.0/legalcode> for more information.



## REFERENCES

1. Mazzini M, Callaini G, Mencarelli C. A comparative analysis of evolution of the egg envelopes and the origin of the yolk. *Ital J Zool.* 1984;51:35-101. doi:[10.1080/1125000849439457](https://doi.org/10.1080/1125000849439457)
2. McMillan DB. Ovulation. In: McMillan DB, ed. *Fish Histology, Female Reproductive Systems*. Springer; 2007:209-284. doi:[10.1007/978-1-4020-5715-1](https://doi.org/10.1007/978-1-4020-5715-1)
3. Beiras R, Bellas J, Cachot J, et al. Ingestion and contact with polyethylene microplastics does not cause acute toxicity on marine zooplankton. *J Hazard Mater.* 2018;360:452-460. doi:[10.1016/j.jhazmat.2018.07.101](https://doi.org/10.1016/j.jhazmat.2018.07.101)
4. Chemello G, Faraoni V, Notarstefano V, Maradonna F, Carnevali O, Gioacchini G. First evidence of microplastics in the yolk and embryos of common cuttlefish (*Sepia officinalis*) from the Central Adriatic Sea: evaluation of embryo and hatching structural integrity and development. *Animals.* 2023;13(1):95. doi:[10.3390/ani13010095](https://doi.org/10.3390/ani13010095). PMID:36611704
5. Plastics Europe. *Plastics-the Facts 2018, an Analysis of European Plastics Production*. Demand and Waste Data. Plastics Europe; 2019.
6. Barnes DKA, Galgani F, Thompson RC, Barlaz M. Accumulation and fragmentation of plastic debris in global environments. *Philos T R Soc B.* 2009;364(1526):1985-1998. doi:[10.1098/rstb.2008.0205](https://doi.org/10.1098/rstb.2008.0205). PMID:19528051
7. Eriksen M, Lebreton LCM, Carson HS, et al. Plastic pollution in the world's oceans: more than 5 trillion plastic pieces weighing over 250,000 tons afloat at sea. *PLoS ONE.* 2014;9(12):0111913. doi:[10.1371/journal.pone.0111913](https://doi.org/10.1371/journal.pone.0111913). PMID:25494041
8. Van Sebille E, Wilcox C, Lebreton L, et al. A global inventory of small floating plastic debris. *Environ Res Lett.* 2015;10(12):124006. doi:[10.1088/1748-9326/10/12/124006](https://doi.org/10.1088/1748-9326/10/12/124006)
9. C  zar A, Echevarria F, Gonz  lez-Gordillo Irigoien X, et al. Plastic debris in the open ocean. *P Natl Acad Sci USA.* 2014;111(28):10239-10244. doi:[10.1073/pnas.1314705111](https://doi.org/10.1073/pnas.1314705111)
10. Wu YQ, Tang ZC. Population composition and migratory distribution of cuttlefish (*Sepiella japonica* de rochebrune) in the Huanghe estuary and the Laizhou gulf. *J Fish China.* 1990;14(2):149-152. doi:[10.1371/journal.ppat.1003386](https://doi.org/10.1371/journal.ppat.1003386). PMID:23825941
11. Chang KM, Wu CW, Lu ZM, Zhu AY, Zhang JS. Comparison in biochemistry of tissues of wild and cultured *Sepiella maindroni*. *Oceanol Limnol Sin.* 2008;39(2):145-151.
12. Wu CW, Dong ZY, Chi CF, Ding F. Reproductive and spawning habits of *Sepiella maindroni* of Zhejiang, China. *Oceanol Limnol Sin.* 2010;41:39-46. doi:[10.11693/hyhz201001005005](https://doi.org/10.11693/hyhz201001005005)
13. Zhao HJ, Xu JK, Yan ZH, Ren HQ, Zhang Y. Microplastics enhance the developmental toxicity of synthetic phenolic antioxidants by disturbing the thyroid function and metabolism in developing zebrafish. *Environment International.* 2020;140:105750. doi:[10.1016/j.envint.2020.105750](https://doi.org/10.1016/j.envint.2020.105750)
14. Yin F, Sun P, Peng SM, et al. The respiration, excretion and biochemical response of the juvenile common Chinese cuttlefish, *Sepiella maindroni* at different temperatures. *Aquaculture.* 2013;402:127-132. doi:[10.1016/j.aquaculture.2013.03.018](https://doi.org/10.1016/j.aquaculture.2013.03.018)
15. Batel A, Borchert F, Reinwald H, Erdinger L, Braunbeck T. Microplastic accumulation patterns and transfer of benzo[a]pyrene to adult zebrafish (*Danio rerio*) gills and zebrafish embryos. *Environm Pollut.* 2018;235:918-930. doi:[10.1016/j.envpol.2018.01.028](https://doi.org/10.1016/j.envpol.2018.01.028)
16. Duan Z, Duan X, Zhao S, et al. Barrier function of zebrafish embryonic chorions against microplastics and nanoplastics and its impact on embryo development. *J Hazard Mater.* 2020;395:122621. doi:[10.1016/j.jhazmat.2020.122621](https://doi.org/10.1016/j.jhazmat.2020.122621)
17. Wan Z, Wang C, Zhou J, et al. Effects of polystyrene microplastics on the composition of the microbiome and metabolism in larval zebrafish. *Chemosphere.* 2019;217:646-658. doi:[10.1016/j.chemosphere.2018.11.070](https://doi.org/10.1016/j.chemosphere.2018.11.070)
18. Duan Y, Xiong D, Wang Y, Li H, Dong H, Zhang J. Toxic effects of ammonia and thermal stress on the intestinal microbiota and transcriptomic and metabolomic responses of *Litopenaeus vannamei*. *Sci Total Environ.* 2021;754:141867. doi:[10.1016/j.scitotenv.2020.141867](https://doi.org/10.1016/j.scitotenv.2020.141867)
19. Yang L, Lv L, Liu H, Wang M, Sui Y, Wang Y. Effects of ocean acidification and microplastics on microflora community composition in the digestive tract of the thick shell Mussel *Mytilus coruscus* through 16S RNA gene sequencing. *B Environ Contam Tox.* 2021;107(4):616-625. doi:[10.1007/s00128-020-03022-5](https://doi.org/10.1007/s00128-020-03022-5)

20. Xu XY, Lee WT, Chan AKY, Lo HS, Shin PKS, Cheung SG. Microplastic ingestion reduces energy intake in the clam *Atactodea striata*. *Mar Pollut Bull.* 2017;124(2):798-802. doi:[10.1016/j.marpolbul.2016.12.027](https://doi.org/10.1016/j.marpolbul.2016.12.027)
21. Paulij WP, Bogaards RH, Denucé JM. Influence of salinity on embryonic development and the distribution of *Sepia officinalis* in the Delta Area (South Western part of The Netherlands). *Mar Biol.* 1990;107(1):17-23. doi:[10.1007/bf01313238](https://doi.org/10.1007/bf01313238)
22. Bolger AM, Lohse M, Usadel B. Trimmomatic: a flexible trimmer for Illumina sequence data. *Bioinformatics.* 2014;30(15):2114-2120. doi:[10.1093/bioinformatics/btu170](https://doi.org/10.1093/bioinformatics/btu170). PMID:24695404
23. Reyon D, Tsai SQ, Khayter D, Foden JA, Sander JD, Joung JK. FLASH assembly of TALENs for high-throughput genome editing. *Nat Biotechnol.* 2012;30(5):460-465. doi:[10.1038/nbt.2170](https://doi.org/10.1038/nbt.2170). PMID:22484455
24. Caporaso JG, Kuczynski J, Stombaugh J, et al. QIIME allows analysis of high-throughput community sequencing data. *Nat Methods.* 2010;7(5):335-336. doi:[10.1038/nmeth.f.303](https://doi.org/10.1038/nmeth.f.303). PMID:20383131
25. Edgar RC, Haas BJ, Clemente JC, Quince C, Knight R. UCHIME improves sensitivity and speed of chimera detection. *Bioinformatics.* 2011;27(16):2194-2200. doi:[10.1093/bioinformatics/btr381](https://doi.org/10.1093/bioinformatics/btr381). PMID:21700674
26. Wang Q, Garrity GM, Tiedje JM, Cole JR. Naive Bayesian classifier for rapid assignment of rRNA sequences into the new bacterial taxonomy. *Appl Environ Microbiol.* 2007;73(16):5261-5267. doi:[10.1128/aem.00062-07](https://doi.org/10.1128/aem.00062-07). PMID:17586664
27. Altschul SF, Gish W, Miller W, Myers EW, Lipman DJ. Basic local alignment search tool. *J Mol Biol.* 1990;215(3):403-410. doi:[10.1016/S0022-2836\(05\)80360-2](https://doi.org/10.1016/S0022-2836(05)80360-2)
28. Lönnstedt OM, Eklöv P. Environmentally relevant concentration of microplastic particles influence larval fish ecology. *Science.* 2016;352(6290):1213-1216. doi:[10.1126/science.aad882](https://doi.org/10.1126/science.aad882)
29. Malafaia G, Souza AMD, Pereira AC, et al. Development toxicity in zebrafish exposed to pristine polyethylene microplastics under static and semi-static systems. *Sci Total Environ.* 2020;700:134867. doi:[10.1016/j.scitotenv.2019.134867](https://doi.org/10.1016/j.scitotenv.2019.134867)
30. Zhao HJ, Xu JK, Yan ZH, Ren HQ, Zhang Y. Microplastics enhance the developmental toxicity of synthetic phenolic antioxidants by disturbing the thyroid function and metabolism in developing zebrafish. *Environment International.* 2020;140:105750. doi:[10.1016/j.envint.2020.105750](https://doi.org/10.1016/j.envint.2020.105750)
31. Bunge A, Kammann U, Scharsack JP. Exposure to microplastic fibers does not change fish early life stage development of three-spined sticklebacks (*Gasterosteus aculeatus*). *Microplastics and Nanoplastics.* 2021;1:15. doi:[10.1186/s43591-021-00015-x](https://doi.org/10.1186/s43591-021-00015-x)
32. Jakubowska M, Białowas M, Stankevičiūtė M, et al. Effects of chronic exposure to microplastics of different polymer types on early life stages of sea trout *Salmo trutta*. *Sci Total Environ.* 2020;740:139922. doi:[10.1016/j.scitotenv.2020.139922](https://doi.org/10.1016/j.scitotenv.2020.139922)
33. Li Y, Wang J, Yang G, et al. Low level of polystyrene microplastics decreases early developmental toxicity of phenanthrene on marine medaka (*Oryzias melastigma*). *J Hazard Mater.* 2020;385:121586. doi:[10.1016/j.jhazmat.2019.121586](https://doi.org/10.1016/j.jhazmat.2019.121586)
34. Jiang W, Fang J, Du M, Gao Y, Fang J, Jiang Z. Microplastics influence physiological processes, growth and reproduction in the Manila clam, *Ruditapes philippinarum*. *Environ Pollut.* 2022;293:118502. doi:[10.1016/j.envpol.2021.118502](https://doi.org/10.1016/j.envpol.2021.118502)
35. Wang Y, Li L, Hu M, Lu W. Physiological energetics of the thick shell mussel *Mytilus coruscus* exposed to seawater acidification and thermal stress. *Sci Total Environ.* 2015;514:261-272. doi:[10.1016/j.scitotenv.2015.01.092](https://doi.org/10.1016/j.scitotenv.2015.01.092)
36. Kong N, Han S, Fu Q, Yu Z, Wang L, Song L. Impact of ocean acidification on the intestinal microflora of the Pacific oyster *Crassostrea gigas*. *Aquaculture.* 2022;546:737365. doi:[10.1016/j.aquaculture.2021.737365](https://doi.org/10.1016/j.aquaculture.2021.737365)
37. Kong N, Zhao J, Zhao B, et al. Effects of high temperature stress on the intestinal histology and microbiota in Yesso scallop *Patinopecten yessoensis*. *Mar Environ Res.* 2023;185:105881. doi:[10.1016/j.marenvres.2023.105881](https://doi.org/10.1016/j.marenvres.2023.105881)
38. Liu H, KarSoon T, Zhang X, et al. Comparison of gut microbiota between golden and brown noble scallop *Chlamys nobilis* and its association with carotenoids. *Front Microbiol.* 2020;11:36. doi:[10.3389/fmicb.2020.00036](https://doi.org/10.3389/fmicb.2020.00036). PMID:32117095
39. Shin NR, Whon TW, Bae JW. Proteobacteria: microbial signature of dysbiosis in gut microbiota. *Trends Biotechnol.* 2015;33(9):496-503. doi:[10.1016/j.tibtech.2015.06.011](https://doi.org/10.1016/j.tibtech.2015.06.011)

40. Brinkmann BW, Beijk WF, Vlieg RC, et al. Adsorption of titanium dioxide nanoparticles onto zebrafish eggs affects colonizing microbiota. *Aquat Toxicol.* 2021;232:105744. doi:[10.1016/j.aquatox.2021.105744](https://doi.org/10.1016/j.aquatox.2021.105744)
41. Hung NV, Schryver PD, Dung NV, Nevejan N, Bossier P. *Ralstonia eutropha*, containing high poly- $\beta$ -hydroxybutyrate levels, regulates the immune response in mussel larvae challenged with *Vibrio coralliilyticus*. *Fish Shellfish Immun.* 2019;84:196-203. doi:[10.1016/j.fsi.2018.09.066](https://doi.org/10.1016/j.fsi.2018.09.066)
42. Brown MR, Barrett SM, Volkman JK, Nearhos SP, Nell JA, Allan GL. Biochemical composition of new yeasts and bacteria evaluated as food for bivalve aquaculture. *Aquaculture.* 1996;143(3-4):341-360. doi:[10.1016/0044-8486\(96\)01286-0](https://doi.org/10.1016/0044-8486(96)01286-0)
43. Jacoby GA. Transmissible antibiotic resistance. In: *Antimicrobial Resistance in the 21st Century*. Springer; 2018:341-381. doi:[10.1007/978-3-319-78538-7\\_25](https://doi.org/10.1007/978-3-319-78538-7_25)
44. Lu L, Luo T, Zhao Y, Cai C, Fu Z, Jin Y. Interaction between microplastics and microorganism as well as gut microbiota: a consideration on environmental animal and human health. *Sci Total Environ.* 2019;667:94-100. doi:[10.1016/j.scitotenv.2019.02.380](https://doi.org/10.1016/j.scitotenv.2019.02.380)